

To Appear in the Astronomical Journal, October 2000

A Search for Stars of Very Low Metal Abundance. V. Photoelectric *UBV* Photometry of Metal-Weak Candidates from the Northern HK Survey

P. Bonifacio¹, S. Monai¹, & T.C. Beers²

ABSTRACT

We report photoelectric *UBV* data for 268 metal-poor candidates chosen from the northern HK objective-prism/interference-filter survey of Beers and colleagues. Over 40 % of the stars have been observed on more than one night, and most have at least several sets of photometric measurements. Reddening estimates, preliminary spectroscopic measurements of abundance, and type classifications are reported.

Subject headings: stars: abundances — stars: Population II — techniques: photometric, spectroscopic

1. Introduction

The HK objective-prism/interference-filter survey, started in 1978 by Preston & Sackett (see Beers, Preston, & Sackett 1985; Beers, Preston, & Sackett 1992), has proven to be a most effective method for the discovery of relatively bright metal-deficient stars in the thick disk and halo of the Galaxy, and is the primary modern source of stars known with $[\text{Fe}/\text{H}] < -3.0$. Such stars are of crucial importance, as they provide fundamental insight on the nucleosynthetic history of the early Galaxy.

The spectroscopic and photometric follow-up of candidate metal-poor stars from the HK survey has become a large international effort, with observational programs being conducted

¹Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, 34131-Trieste, Italy;
email: bonifaci@ts.astro.it, monai@ts.astro.it

²Michigan State University, Department of Physics & Astronomy, E. Lansing, MI 48824;
email: beers@pa.msu.edu

by colleagues with access to both northern and southern hemisphere telescopes. Besides the identification of extremely metal-poor stars, the HK survey is providing a large number of stars of intermediate metallicity ($-0.5 \leq [\text{Fe}/\text{H}] \leq -2.0$) – stars that are useful for the study of the thick disk, the metal-weak thick disk (MWTD), and of the thick disk to halo transition.

Since the HK survey is kinematically unbiased in its selection criteria, it is also an ideal resource for studies of the kinematics and dynamics of the halo and thick-disk populations. Many of the brighter HK survey candidates have available proper motion measurements, and, when combined with accurate (photometric, at present) distance estimates and radial velocities, yield a large sample of stars with full space motions (e.g., Beers et al. 2000). Based on such detailed information, Chiba & Beers (2000) explore a number of important issues related to the formation and evolution of the Galaxy, and show how the kinematics of local samples can be used to provide constraints on the low-metallicity cutoff of the MWTD population ($[\text{Fe}/\text{H}] \sim -2.0$), the degree of flattening in the thick-disk and halo populations, and the variation of flattening with Galactocentric distance, the existence (or not) of the supposed “counter-rotating high-halo population,” measurements of velocity gradients in the thick disk and halo, and searches for kinematic substructure such as reported by Helmi et al. (1999).

A large campaign to obtain medium-resolution spectroscopic observations of additional HK survey metal-poor candidates has been underway for roughly the past seven years, and the total number of stars with suitable spectra for abundance and radial velocity measurement is now on the order of $N \sim 5000$ (see Beers 1999 for a summary). Unfortunately, the photometric follow-up campaign has been lagging behind somewhat, especially in the northern hemisphere, due to the lack of suitable telescopes (and sites) for carrying out the observations (a problem that is getting worse, not better, as time passes). The bulk of the extant northern *UBV* data comes from a study of 300 HK candidates by Doinidis & Beers (1990, 1991). Strömgren photometry of a sample of over 500 northern HK survey stars has been recently reported by Anthony-Twarog et al. (2000). Additional Strömgren photometry of 140 northern HK metal-poor candidates has been obtained by Schuster et al. (2000). In the southern hemisphere the situation is somewhat better, though additional work is still required. Preston, Shectman, & Beers (1991) obtained *UBV* for a sample of ~ 1800 southern HK survey stars, though most were field horizontal-branch (hereafter, FHB) stars rather than cooler metal-poor candidates. *UBV* measurements for a large sample ($N \sim 2600$) of southern HK survey candidate metal-poor stars have been recently reported by Norris, Ryan, & Beers (1999). Schuster et al. (1996) report Strömgren photometry for a small ($N \sim 90$) sample of metal-poor HK survey stars, including both northern and southern candidates.

The photometry is important mainly for two reasons – (1) to improve the metallicity estimates of the stars, making use of the de-reddened $(B - V)_0$ color (or its approximation from the Strömgren $(b - y)_0$ index) and the calibration procedure described by Beers et al. (1999), and (2) for obtaining photometric distance estimates. For the latter purpose it is important to also have the $(U - B)_0$ color, since it enables one to discriminate (in most, but not all cases) between FHB stars and main-sequence gravity stars near the turnoff of an old stellar population (hereafter, TO) in the color range $0.3 \leq (B - V)_0 \leq 0.5$.

We thus decided to initiate a northern photometric campaign with a small (91 cm) telescope (somewhat undersized for photoelectric follow-up of our targets, which have apparent magnitudes mainly in the range $12.5 < V < 14.5$) for which we were granted substantial amounts of observing time over an extended period. The results of the first year of this campaign (from August 1998 to August 1999) are quite satisfactory. We obtained data on 22 nights out of 30 allocated, and here we report UBV for a total of 268 stars. Roughly 40 % of these were observed on more than one night. For most stars observations on different nights were consistent and their average provides accurate magnitudes and colors, but for a small number of stars they were not. We also present data for these stars, flagging them as “suspected variables.” All but a handful of our stars already have medium-resolution spectroscopy available, so the reported photometry can be put to immediate use.

2. Observations and Data Reduction

The observations were carried out with the 91 cm telescope of the Osservatorio Astrofisico di Catania at the M.G. Fracastoro mountain station on Mt. Etna during 4 runs in August 1998, October 1998, May 1999, and August 1999. A log of the observations is given in Table 1, along with estimates of the typical residuals (relative to the photometric standard stars) that were achieved over the course of the run. All observations were performed with a single-channel photoelectric photometer mounted at the Cassegrain focus. The photomultiplier tube was a EMI 9893QA/350, which mimics a S20 spectral response, used along with Johnson UBV filters. For all the observations we employed diaphragm #2, corresponding to a $21''.7$ diameter on the sky. Calibrations were obtained by observations of typically 10–15 standard stars each night, taken from the lists of Landolt (1983, 1992). The calibration scheme followed the method outlined by Harris, Fitzgerald, & Reed (1981), retaining only linear terms in the colors. This method allows one to tie several different nights into a single calibration, and hence recover usable data for “partial” nights during which few standard stars were observed due to changes in the weather conditions.

Given its location at the foot of an active volcano, the Serra La Nave Observatory is in

a somewhat particular situation. Typically the wind blows from west to east, thus keeping smoke from the volcano away from the Observatory. When the wind blows from east to west the smoke moves in the direction of the Observatory, and it is necessary to close the dome to avoid the dust damaging the mirror. In addition, there are many intermediate situations in which, either due to a slightly different direction of the wind or to extra activity of the volcano, the measured extinction is markedly different from the “typical” values – these nights may well be photometric, however they must be calibrated on their own and may not be tied to other nights.

Our reduction scheme began by performing a calibration of each night individually, enabling rejection of “bad” standards (usually easily identified as outliers). Next we compared the derived extinctions of the different nights. Nights with highly deviant extinctions were left with their individual calibration, while the others were tied together with a single “multi-night” calibration. On a few “odd” nights it was necessary to introduce a linear term in UT into the extinction equations. In Table 1 we indicate the nights that have been treated with the “multi-night” reduction scheme (indicated by an “m” in the fifth column), and the ones that were calibrated individually (indicated by an “s” in the fifth column). A “t” in the last column denotes nights for which a time-dependent extinction has been assumed. No attempt was made to tie observations taken on different runs into a single calibration, because the seasonal change in the extinction is likely to add uncertainty to the resulting calibration.

Once all the data had been calibrated we averaged together the available observations of each star. Some stars were observed on only one night, even though several times, while others were observed on several different nights. These stars broadly divided into two groups: stars whose standard deviations in the derived magnitudes were less than or equal to the expected error and others for which they were larger. We define the first group as “stars that display no variability,” and list their photometric information in Table 2. Column (1) of this table lists the star name. Columns (2) and (3) are the (1950) epoch coordinates (accurate to several arc-seconds), and columns (4) and (5) are the (2000) epoch coordinates. Columns (6) and (7) are the corresponding Galactic coordinates of each star. Columns (8)-(13) list the measured V , $B - V$, and $U - B$, and their associated one-sigma errors. Columns (14) and (15) list the numbers of independent nights on which the star was observed, and the total number of (useful) observations per star that were employed, respectively. The final two columns of this table list two alternative estimates of reddening – column (16) is that derived from the new reddening maps of Schlegel, Finkbeiner, & Davis (1998), while column (17) is that obtained from the maps of Burstein & Heiles (1982). We describe our choice of reddening below.

The second set of stars are reported as “suspected variables.” As a dividing line we

chose the standard deviation in the V magnitude: if $\sigma_V > 0.08$ mag we consider the star to be a “suspected variable.” This division is to some extent arbitrary and the status of “suspected variables” should mainly be used as a warning flag. For all the “suspected variables” that were observed on at least three different nights we computed a Lomb normalized periodogram, using the routine `period` of Press et al (1992). Only the star BS 16542–0002 displayed peaks with a significance above 95% – we therefore consider this as the only confirmed variable of our present sample. This star exhibits colors $((B - V)_0 = 0.70)$ that exclude it from being an RR Lyrae variable, so it merits additional attention to establish the source of its apparent variation. The periodogram analysis is optimized for periodic variables, so that any irregular or semi-regular variables are likely to go undetected by this criterion. In Table 3, we list the mean magnitudes and colors, and their standard deviations, for “suspected” variables, following the same column assignments as in Table 2.

Note that a number of objective-prism plates in the HK survey overlap with one another, hence there are several instances of candidates with multiple names. Table 4 summarizes the multiple identifications for the HK survey stars in our present sample.

3. External Comparisons

There are 87 stars in common between the present paper and the paper of Anthony-Twarog et al. (2000), which provides *uvbyCa* photometry. Our V magnitudes may be directly compared to theirs, which were obtained by calibrating the y filter to the Johnson V system. In two cases the difference is large enough that we may be almost certain the we did not observe the same star. In the case of BS 17570–0057 our measurements indicate $V = 13.40$, $B - V = 0.43$, while Anthony-Twarog et al. give $V = 9.972$, $B - V \approx 1.35 \times b - y = 0.87$. The Guide Star Catalog lists an estimated V magnitude for this star of $V = 13.4$, and our medium-resolution spectra provide an estimate of $B - V \sim 0.47$, based on the calibration of the HP2 Balmer-line index described in Beers et al. (1999). We observed the star on two different nights for a total of four measurements, and the measurements were all consistent with one another. In the case of BS 17583–0012 the situation is less clear. We observed the star on one night only, and obtain $V = 12.45$, $B - V = 0.54$, while Anthony-Twarog et al. give $V = 11.497$, and from their reported $b - y$ color we estimate $B - V \approx 0.83$. For this star the Guide Star Catalog lists $V = 12.1$, and from the HP2 index we estimate $B - V \sim 0.39$. Hence it is possible that we observed different stars, possibly neither of which was the correct one, however it is also possible that the star is variable. We exclude these two stars from the comparison. Future measurements will permit us to establish the correct photometry for both stars and to check for variability.

The top panel of Figure 1 shows a comparison of the V magnitudes reported by Anthony-Twarog et al., V (AT), with our measurements for the remaining 85 stars—the agreement is quite satisfactory. We estimate the difference in central location (“mean”) between these two samples, and the scale (“standard deviation”) of their differences, using the robust and resistant biweight estimators described by Beers, Flynn, & Gebhardt (1990). We obtain $C_{BI}(\Delta V) = -0.003$ magnitudes, and $S_{BI}(\Delta V) = 0.053$ magnitudes. The biweight scale may be considered as a conservative estimate of the external accuracy of our photometry, if we assume that the V magnitudes of Anthony-Twarog et al. are error-free, and the scatter in ΔV is due solely to the errors in our photometry. A more realistic error estimate is obtained by assuming that the external errors of the photometry of both groups is the same and dividing the above value by $\sqrt{2}$, obtaining 0.04 magnitudes, which is on the order of the reliability of the reddening corrections discussed below.

The lower panel of Figure 1 shows the comparison of the estimated $B - V$ color of Anthony-Twarog et al., derived from the standard relation $B - V_{est} = 1.35 \times b - y$, with our measured $B - V$ colors. The agreement is generally good, with the possible exception of the bluest stars, which could of course include bona-fide variables such as RR Lyraes. The biweight estimate of central location of the difference in color $\Delta(B - V) = (B - V)_{(our)} - (B - V)_{est}(AT)$ is $C_{BI} = -0.02$ magnitudes. The biweight scale estimate of the difference in colors is $S_{BI} = 0.06$ magnitudes. Apportioning the errors equally between the two studies suggests an external accuracy of our $B - V$ values of ~ 0.04 magnitudes. This is perhaps a bit too pessimistic, since we are comparing two different photometric systems (Strömgren versus Johnson), and have assumed that the transformation equations are both error free and apply equally well over the range of colors and abundances of our program stars.

4. Adopted Reddenings, Preliminary Abundances, and Stellar Classifications

We now consider the nature of our target stars, following procedures described by Beers et al. (1999), slightly modified as described below.

4.1. Reddening Estimates

We initially adopted the Schlegel et al. (1998) estimates of reddening from Tables 2 and 3, which have superior spatial resolution and are thought to have a better-determined zero point than the Burstein & Heiles (1982) maps. However, Arce & Goodman (1999) caution that the Schlegel et al. map may overestimate the reddening values when the color

excess $E (B - V)_S$ exceeds about 0.15 magnitudes. Our own independent tests suggest that this problem may extend to even lower color excesses, on the order of $E (B - V)_S = 0.10$ magnitudes. Hence, we have adopted a slight revision of the Schlegel et al. reddening estimates, according to the following:

$$\begin{aligned} E (B - V)_A &= E (B - V)_S & E (B - V)_S &\leq 0.10 \\ E (B - V)_A &= 0.10 + 0.65 \times [E (B - V)_S - 0.10] & E (B - V)_S &> 0.10 \end{aligned} \tag{1}$$

where $E (B - V)_A$ indicates the adopted reddening estimate. We note that for $E (B - V)_S \geq 0.15$ this approximately reproduces the 30%–50% reddening reduction recommended by Arce & Goodman. To account for stars that are located within the reddening layer, assumed to have a scale height $h = 125$ pc, the reddening to a given star at distance D is reduced compared to the total reddening by a factor $[1 - \exp(-|D \sin b|/h)]$, where b is the Galactic latitude. The distance is estimated from M_V vs. $(B - V)_0$ relations, as described in Beers et al. (2000). The procedure must be iterated, because both V_0 (and therefore D) and $(B - V)_0$ depend on the $E (B - V)_A$. Since the M_V vs. $(B - V)_0$ relation depends on metallicity, as well as on the classification of the star, at each step of the iteration the metallicity is recomputed and the classifications redetermined with the current estimates of $(B - V)_0$ and $(U - B)_0$, so that at the end we obtain consistent estimates of $E (B - V)_A$, $[\text{Fe}/\text{H}]$, and D . The procedure is not without shortcomings, as in some regions of the color and abundance space there is little or no power in discrimination of type classifications. This classification degeneracy adds some uncertainty, as discussed below.

The final adopted estimate of reddening is listed in column (2) of Tables 5 and 6. Column (3) lists the resulting extinction-corrected V_0 magnitudes, using $A_V = 3.1 E (B - V)_A$. Reddening-corrected colors, obtained from $(B - V)_0 = B - V - E (B - V)_A$, and $(U - B)_0 = U - B - 0.72 E (B - V)_A$ are listed in columns (4) and (5), respectively.

4.2. Abundance Estimates

Medium-resolution spectroscopy was obtained for the majority of our program stars with either the 2.1m telescope at Kitt Peak National Observatory, using the Goldcam spectrograph, or the 2.5m Isaac Newton Telescope on La Palma, using the Intermediate Dispersion Spectrograph. For a few stars we have combined estimates obtained from other spectroscopic follow-up campaigns. Details of the acquisition of this data are discussed in Beers et al. (1999) and Allende Prieto et al. (2000), and will not be repeated here. The full set of

medium-resolution spectroscopic results for HK survey stars from the recent campaigns will be presented in subsequent papers.

Measurements of the KP index (a pseudo-equivalent width of the CaII K line) obtained from the spectra were used to obtain the metallicity estimate $[\text{Fe}/\text{H}]_{\text{K3}}$, following the procedures outlined in Beers et al., employing the de-reddened $(B - V)_0$ colors described above. The resulting abundance estimates, and their associated one-sigma errors, are listed in column (6) and (7) of Tables 5 and 6, respectively. For a few of the redder stars, inspection of the spectra makes it clear that the CaII K line exhibits the presence of emission in its core, likely compromising the KP index, and hence, the estimated metallicity. We have refrained from quoting abundances, or making type classifications, for these stars, and have indicated them with the designation “CORE EM” in column (9) of the tables.

For each star with an available spectrum we have also made independent estimates of the expected $(B - V)_0$ color, derived from the calibration of this color with the HP2 Balmer-line index described by Beers et al (1999). In a number of cases, this calibration yielded predicted de-reddened colors that differed from our measured colors by larger than 0.15 magnitudes (2–3 times the expected error in the HP2 calibration) – the abundances of these stars are indicated as somewhat uncertain by an appended “:” in column (6) of Tables 5 and 6. Stars classified as TO/FHB or FHB/A in column (8) of Tables 5 and 6 (as discussed below) also have some uncertainty in their adopted metallicities, which we indicate by an appended “:” to their abundance estimates. A few stars in our present sample have colors that extend beyond the nominal region of the Beers et al. metallicity calibration ($0.30 \leq (B - V)_0 \leq 1.2$). These stars are indicated in Column (9) of the tables with the designation “BLUE” or “RED”, and their abundance estimates have an appended “::” to indicate the additional uncertainty. For the two stars with colors blueward of $(B - V)_0 = 0.0$ no abundance estimates is given.

4.3. Stellar Classifications

Classification of the program stars follows the procedures described in Beers et al. (1999), with the following caveat. The original classification scheme (dating back to Beers et al. 1985) was designed for application to stars of very low metallicity ($[\text{Fe}/\text{H}] < -2.0$). However, since the HK follow-up now includes large numbers of stars with higher abundances, the procedure by which the “split” between FHB and TO stars is made has to be modified. In the original scheme, stars in the color range $0.3 \leq (B - V)_0 \leq 0.5$, which includes both FHB and TO stars, were assigned classifications based on the simple criteria:

$$TO : \quad (U - B)_0 \leq -0.1$$

$$FHB : \quad (U - B)_0 > -0.1$$

This simple criteria is not adequate for stars with abundances $[\text{Fe}/\text{H}] > -2.0$.

Figure 2 shows the “limiting” $(U - B)_0$ colors as a function of metallicity, $[\text{Fe}/\text{H}]$, as predicted from the Revised Yale Isochrones (Green 1988; King, Demarque, & Green 1988) for (a) stars with surface gravities $\log g \leq 3.0$ (appropriate for FHB stars in this color range—see Wilhelm, Beers, & Gray 1999) and (b) stars with $\log g \geq 4.0$ (appropriate for TO stars in this color range). For the present exercise, we make the classification assignments by comparison with these limiting lines (broadened slightly to include possible reddening errors on the order of 0.03 magnitudes). Fits to these lines are well produced by the following quadratic relations:

$$TO : \quad (U - B)_0(\text{lim}) = 0.080 + 0.207 [\text{Fe}/\text{H}] + 0.038 [\text{Fe}/\text{H}]^2 + 0.02$$

$$FHB : \quad (U - B)_0(\text{lim}) = 0.105 + 0.130 [\text{Fe}/\text{H}] + 0.020 [\text{Fe}/\text{H}]^2 - 0.02$$

The 0.02 constant that is added or subtracted in the above relations is the correction term for the reddening error in $(U - B)_0$. Stars with $(U - B)_0$ colors *above* the TO line (i.e., at lower values of $(U - B)_0$) are classified as TO, while those *below* the FHB line (i.e., at higher values of $(U - B)_0$) are classified as FHB. Stars of intermediate surface gravities fall between the lines (as do some stars as a result of errors in their metallicities or colors), and their classifications are less certain. Since we expect MOST of the stars in this color regime to be metal-rich TO stars rather than metal-poor FHB stars, for stars between the two limiting lines we assign the classification TO/FHB (indicating that TO is our preference in the classification, and is the one used for distance determination in the assignment of reddening). In most cases accurate photometry should produce a reliable assignment, which is listed (along with classifications for other types carried out following the Beers et al. 1999 procedures) in column (8) of Tables 5 and 6. Note that for stars with colors blueward of $(B - V)_0 = 0.30$ we have adopted a classification of FHB/A, and do not attempt to refine this assignment further at present (however, see Wilhelm et al. 1999 for a description of how this might be done). For the two stars with colors blueward of $(B - V)_0 = 0.0$ no classification is given.

5. Discussion

Figure 3 is a two-color diagram, $(U - B)_0$ *vs.* $(B - V)_0$, for the stars in our present sample not listed as suspected variables, over the color range for which reliable abundance estimates could be obtained. A solar-abundance main-sequence line, estimated from the Revised Yale Isochrones, is shown for comparison.

All of our program stars have available radial velocities (accurate to $\sim 7 - 10$ km s⁻¹), and many of the brighter ones have proper motions available from a number of sources. We intend to expand our photometric program of the HK survey stars (especially those with available proper motions) and thereby supplement the basic data with which we can carry out a detailed investigation of the kinematics of thick disk and halo stars.

We are grateful to Prof. S. Catalano for generously allocating time to such a long-term project. Special thanks are due to G. Carbonaro, A. Di Stefano, G. Occhipinti, M. Puleo, and S. Sciuto for their assistance during the observations. We wish to thank E. Marilli and C. Lo Presti for providing the source code of the PHOT reduction package with which we implemented the multi-night reduction scheme. Finally we thank T. Valente for carefully preparing the many finding charts needed for the observations. This work was supported in part by NATO grant 950875. T.C.B also acknowledges partial support from grant AST 95-29454 from the National Science Foundation.

REFERENCES

- Allende Prieto, C., Rebolo, R., Garcia Lopez, R., Serra-Ricart, M., Beers, T.C., Rossi, S., Bonifacio, P., & Molaro, P. 2000, *AJ*, in press. astro-ph/0005598
- Anthony-Twarog, B.J., Sarajedini, A., Twarog, B.A., & Beers, T.C. 2000, *AJ*, in press. astro-ph/0003115
- Arce, H.G., & Goodman, A.A. 1999, *ApJ*, 512, L135
- Beers, T.C. 1999, in *The Third Stromlo Symposium: The Galactic Halo*, eds. B.K. Gibson, T.S. Axelrod, & M.E. Putman, (San Francisco: ASP), Vol. 165, p. 206
- Beers, T.C., Flynn, C., & Gebhardt, K. 1990, *AJ*, 100, 32
- Beers, T.C., Preston, G.W., & Shectman, S.A. 1985, *AJ*, 90, 2089
- Beers, T.C., Preston, G.W., & Shectman, S.A. 1992, *AJ*, 103, 1987
- Beers, T.C., Rossi, S., Norris, J.E., Ryan, S.G., & Shefler, T. 1999, *AJ*, 117, 981.
- Beers, T.C., Chiba, M., Yoshii, Y., Platais, I., Hanson, R.B., Fuchs, B., & Rossi, S. 2000, *AJ*, in press, astro-ph/0003103
- Burstein, D., & Heiles, C. 1982, *AJ*, 87, 1165
- Carney, B.W. 1999, in *The Third Stromlo Symposium: The Galactic Halo*, eds. B.K. Gibson, T.S. Axelrod, & M.E. Putman, (San Francisco: ASP), Vol. 165, p. 230
- Chiba, M., & Beers, T.C. 2000, *AJ*, in press, astro-ph/0003087
- Doinidis, S.P., & Beers, T.C. 1990, *PASP*, 102, 1392
- Doinidis, S.P., & Beers, T.C. 1991, *PASP*, 103, 973
- Green, E.M. 1988, in *Calibration of Stellar Ages*, ed. A.G.D. Philip, (Schenectady: L. Davis Press), p. 81
- Harris, W.E., Fitzgerald, M.P., & Reed, B.C., 1981, *AJ*, 93, 507
- Helmi, A., White, S.D.M., deZeeuw, P.T., & Zhao, H. 1999, *Nature*, 402, 53
- King, C.C., Demarque, P., & Green, E.M. 1988,
- Landolt, A. U., 1983, *AJ*, 88, 439

- Landolt, A. U., 1992, AJ, 104, 340
- Norris, J.E., Ryan, S.G., & Beers, T.C. 1999, ApJS, 123, 639
- Press W.H., Teukolsky S.A., Vetterling W.T., Flannery B.P., 1992 “Numerical Recipes”, Cambridge University Press, Cambridge UK
- Preston, G.W., Sheckman, S.A., & Beers, T.C. 1991, ApJS, 76, 1001
- Schlegel, D.J., Finkbeiner, D.P., & Davis, M. 1998, ApJ, 500, 525
- Schuster, W.J., Nissen, P.E., Parrao, L., Beers, T.C., & Overgaard, L.P. 1996, A&AS, 117, 317
- Schuster, W.J., Parrao, L., Franco, A., Beers, T.C., & Nissen, P.E. 2000, in Galaxy Evolution – Connecting the Distant Universe with the Local Fossil Record, ed. M. Spite (Dordrecht: Kluwer), p. 141
- Wilhelm, R., Beers, T.C., & Gray, R. 1999, AJ, 117, 2308

Table 1. Log of Observations

Date	σ_V	σ_{B-V}	σ_{U-B}	r ^a	Comment
27/08/1998	0.012	0.015	0.013	s	
28/08/1998	0.013	0.007	0.012	s	
15/10/1998				m	
16/10/1998				m	
18/10/1998				m	
21/10/1998	0.008	0.008	0.010	m	
22/10/1998	0.024	0.012	0.010	s	
13/05/1999				m	
15/05/1999				m	
16/05/1999				m	
18/05/1999				m	
20/05/1999	0.028	0.014	0.018	m	
5/08/1999				m	
6/08/1999				m	
8/08/1999				m	
11/08/1999				m	
13/08/1999				m	
14/08/1999				m	
15/08/1999	0.017	0.013	0.022	m	
7/08/1999	0.031	0.012	0.014	s	
9/08/1999	0.015	0.015	0.028	s	t
10/08/1999	0.010	0.015	0.021	s	t

^areduction type: m=multi-night; s=single-night

Table 2. Photometric Data For Program Stars

Star	RA	DEC	RA	DEC	l	b	V	σ_V	$B - V$	σ_{B-V}	$U - B$	σ_{U-B}	N	N	$E(B - V)$	$E(B - V)$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14) ⁿⁱ	(15) ^{ob}	(16) ^S	(17) ^B
BS 15622-0053	12 51 07.7	+25 21 58	12 53 34.3	+25 05 42	316.3	87.9	13.59	0.01	0.74	0.03	0.38	0.06	1	2	0.01	0.02
BS 15622-0055	12 48 55.6	+27 11 30	12 51 21.9	+26 55 12	298.5	89.8	14.05	0.03	0.40	0.02	-0.17	0.01	1	2	0.01	0.00
BS 15624-0040	16 15 27.3	+46 11 08	16 16 58.5	+46 03 49	72.2	45.5	13.57	0.03	0.41	0.02	-0.22	0.02	3	6	0.01	0.00
BS 16034-0101	15 48 50.0	+54 49 36	15 50 04.1	+54 40 35	85.9	47.6	12.36	0.01	0.43	0.01	-0.04	0.01	1	2	0.01	0.00
BS 16034-0115	15 59 04.4	+57 26 38	16 00 07.9	+57 18 15	88.7	45.3	11.98	0.02	0.37	0.01	-0.26	0.01	2	4	0.02	0.00
BS 16076-0006	12 45 54.2	+21 13 02	12 48 22.6	+20 56 41	296.3	83.8	13.36	0.01	0.52	0.02	-0.19	0.02	1	4	0.03	0.03
BS 16076-0012	12 49 25.2	+18 02 29	12 51 54.2	+17 46 11	303.6	80.6	13.76	0.05	0.47	0.02	-0.26	0.03	1	2	0.02	0.01
BS 16076-0040	12 54 47.9	+22 49 28	12 57 14.8	+22 33 16	319.3	85.2	12.00	0.01	0.40	0.01	-0.27	0.03	1	2	0.04	0.03
BS 16076-0050	12 55 49.0	+18 11 45	12 58 17.3	+17 55 34	313.0	80.7	13.11	0.03	0.56	0.03	-0.25	0.01	1	2	0.04	0.03
BS 16077-0007	11 32 39.7	+31 17 05	11 35 18.0	+31 00 29	195.3	72.9	12.21	0.02	0.33	0.02	-0.28	0.01	1	2	0.02	0.00
BS 16077-0023	11 37 29.0	+31 03 59	11 40 06.4	+30 47 21	195.5	74.0	12.24	0.01	0.37	0.02	-0.27	0.01	1	2	0.02	0.00
BS 16077-0047	11 44 27.6	+32 02 22	11 47 03.9	+31 45 42	190.8	75.3	14.40	0.01	0.53	0.06	-0.30	0.09	1	2	0.02	0.01
BS 16077-0077	11 52 41.9	+28 37 44	11 55 16.5	+28 21 02	204.1	77.5	13.70	0.04	0.61	0.01	-0.14	0.07	1	2	0.03	0.00
BS 16078-0025	13 33 20.1	+31 17 46	13 35 37.6	+31 02 28	57.6	79.6	12.61	0.01	0.42	0.01	-0.21	0.01	1	2	0.02	0.00
BS 16079-0019	15 45 40.8	+60 14 41	15 46 37.4	+60 05 28	93.3	45.7	14.09	0.01	0.38	0.01	-0.23	0.01	1	4	0.02	0.01
BS 16079-0133	16 14 13.5	+57 19 27	16 15 14.1	+57 12 02	87.6	43.4	12.06	0.01	0.43	0.01	0.00	0.01	1	6	0.01	0.00
BS 16080-0007	16 27 45.1	+59 33 17	16 28 34.4	+59 26 46	89.7	41.1	13.46	0.01	0.46	0.01	-0.22	0.03	2	4	0.01	0.00
BS 16080-0022	16 30 59.8	+59 37 44	16 31 48.2	+59 31 26	89.7	40.6	13.13	0.01	0.43	0.01	-0.22	0.01	2	4	0.02	0.00
BS 16080-0054	16 38 37.9	+61 03 29	16 39 18.5	+60 57 42	91.2	39.4	12.79	0.02	0.71	0.01	0.20	0.03	3	6	0.03	0.00
BS 16080-0067	16 42 31.5	+58 46 08	16 43 21.4	+58 40 37	88.1	39.4	13.15	0.02	0.54	0.01	-0.06	0.01	2	4	0.01	0.00
BS 16080-0093	16 49 51.9	+59 18 06	16 50 38.5	+59 13 05	88.6	38.4	13.61	0.01	0.69	0.01	0.13	0.05	4	9	0.02	0.00
BS 16080-0175	16 50 14.3	+57 17 23	16 51 08.9	+57 12 24	86.0	38.7	14.35	0.03	0.45	0.02	-0.16	0.02	2	4	0.03	0.02
BS 16082-0029	13 47 29.8	+28 32 53	13 49 47.0	+28 18 01	41.5	77.0	13.17	0.02	0.59	0.01	0.06	0.01	3	6	0.01	0.00
BS 16082-0045	13 53 15.0	+31 56 43	13 55 28.9	+31 42 03	54.9	75.3	13.31	0.02	0.40	0.01	-0.26	0.02	3	7	0.02	0.00
BS 16082-0097	14 02 18.8	+32 37 02	14 04 30.7	+32 22 42	55.7	73.3	12.12	0.01	0.36	0.01	-0.18	0.01	1	2	0.02	0.00
BS 16082-0129	13 44 54.4	+29 12 43	13 47 11.5	+28 57 46	44.7	77.6	13.61	0.02	0.62	0.02	0.04	0.01	2	5	0.02	0.00
BS 16083-0020	14 46 44.1	+49 13 19	14 48 26.3	+49 00 53	85.0	58.6	13.64	0.01	0.31	0.01	-0.21	0.02	1	2	0.02	0.00
BS 16083-0024	14 45 52.2	+50 41 39	14 47 31.9	+50 29 10	87.4	57.9	14.34	0.02	0.48	0.04	-0.07	0.08	2	8	0.04	0.01
BS 16083-0043	14 51 29.3	+47 15 00	14 53 13.8	+47 02 48	80.9	58.9	13.16	0.01	0.39	0.01	-0.29	0.01	2	6	0.02	0.00
BS 16083-0065	14 56 25.2	+51 52 24	14 57 59.7	+51 40 27	87.5	55.9	14.80	0.03	0.44	0.03	-0.18	0.06	2	4	0.02	0.00
BS 16083-0146	15 06 54.2	+47 26 10	15 08 35.0	+47 14 46	79.1	56.5	14.80	0.03	0.28	0.03	-0.29	0.05	2	6	0.03	0.01
BS 16084-0123	16 21 53.7	+54 30 19	16 23 02.3	+54 23 25	83.4	43.1	13.56	0.01	0.42	0.01	-0.01	0.02	1	2	0.01	0.00
BS 16084-0160	16 27 43.8	+54 43 33	16 28 50.7	+54 37 02	83.5	42.2	13.12	0.01	0.82	0.01	0.28	0.01	2	4	0.01	0.00
BS 16089-0013	13 47 10.4	+35 59 45	13 49 22.1	+35 44 53	71.5	75.0	13.33	0.02	0.70	0.01	0.08	0.02	3	8	0.01	0.00
BS 16089-0042	13 51 26.6	+35 01 18	13 53 38.3	+34 46 34	66.6	74.7	14.20	0.01	0.40	0.01	-0.01	0.02	1	2	0.02	0.00
BS 16089-0086	14 00 16.3	+34 32 06	14 02 26.9	+34 17 41	62.5	73.2	14.11	0.03	0.32	0.01	0.10	0.03	1	2	0.02	0.00
BS 16468-0029	09 03 04.1	+41 34 35	09 06 18.7	+41 22 32	180.4	42.2	13.54	0.01	0.41	0.02	-0.27	0.03	1	4	0.02	0.00
BS 16542-0016	16 20 51.5	+20 22 15	16 23 02.7	+20 15 19	36.8	41.4	13.56	0.01	0.73	0.01	-0.32	0.04	1	3	0.06	0.06
BS 16542-0022	16 22 15.5	+21 47 02	16 24 24.9	+21 40 11	38.7	41.6	12.94	0.01	0.43	0.02	-0.28	0.01	2	8	0.05	0.03
BS 16542-0051	16 25 55.4	+20 50 37	16 28 05.8	+20 44 01	37.9	40.4	14.52	0.02	0.52	0.02	-0.04	0.03	2	5	0.05	0.05
BS 16542-0052	16 25 33.7	+21 10 29	16 27 43.7	+21 03 52	38.2	40.6	12.76	0.01	0.87	0.01	0.42	0.01	2	5	0.05	0.05
BS 16545-0074	11 20 55.4	+35 34 45	11 23 36.9	+35 18 16	183.6	69.6	14.09	0.03	0.43	0.04	-0.27	0.03	1	3	0.02	0.01
BS 16545-0082	11 22 38.0	+35 35 16	11 25 19.1	+35 18 46	183.6	69.9	13.02	0.01	0.49	0.02	-0.21	0.02	1	2	0.02	0.00
BS 16545-0089	11 21 45.9	+37 06 57	11 24 27.7	+36 50 28	179.6	69.2	14.38	0.01	0.38	0.02	-0.30	0.01	1	3	0.02	0.01
BS 16548-0028	16 25 12.1	+52 49 51	16 26 25.2	+52 43 10	81.1	43.0	14.65	0.01	0.43	0.05	-0.19	0.01	2	4	0.03	0.00
BS 16548-0070	16 39 31.4	+48 22 55	16 40 54.6	+48 17 13	74.7	41.3	13.69	0.01	0.67	0.02	-0.06	0.05	2	4	0.02	0.00
BS 16548-0140	16 50 24.5	+52 16 15	16 51 35.9	+52 11 18	79.6	39.3	12.61	0.01	0.47	0.02	-0.19	0.01	2	4	0.03	0.00
BS 16549-0017	11 50 05.7	+34 04 11	11 52 41.1	+33 47 30	181.6	75.7	12.42	0.01	0.39	0.01	-0.25	0.01	1	2	0.02	0.01
BS 16552-0038	16 50 53.0	+35 21 51	16 52 41.5	+35 16 57	57.9	38.5	14.60	0.02	0.54	0.02	-0.12	0.01	2	6	0.02	0.01
BS 16552-0042	16 49 35.2	+34 42 57	16 51 24.8	+34 37 57	57.0	38.7	14.64	0.01	0.39	0.01	0.04	0.01	1	2	0.02	0.01
BS 16552-0086	16 58 10.7	+33 35 01	17 00 01.7	+33 30 37	56.0	36.8	14.60	0.03	0.62	0.03	0.15	0.01	1	2	0.02	0.01
BS 16552-0090	16 56 16.8	+34 23 29	16 58 06.6	+34 18 57	56.9	37.3	13.28	0.01	0.74	0.01	0.18	0.02	1	2	0.02	0.01
BS 16552-0142	17 05 17.2	+34 25 29	17 07 06.4	+34 21 36	57.4	35.5	12.05	0.01	0.46	0.02	-0.26	0.01	2	6	0.03	0.02
BS 16554-0053	14 00 42.8	+21 00 46	14 03 03.6	+20 46 22	15.5	72.5	15.05	0.01	0.29	0.05	-0.26	0.06	1	2	0.03	0.01
BS 16556-0045	16 38 18.9	+32 16 26	16 40 13.1	+32 10 40	53.4	40.6	13.94	0.01	0.64	0.01	0.02	0.02	1	2	0.03	0.01
BS 16556-0046	16 39 35.2	+32 11 27	16 41 29.4	+32 05 46	53.3	40.3	14.18	0.01	0.56	0.01	-0.10	0.05	1	2	0.02	0.01
BS 16556-0054	16 40 02.8	+30 37 47	16 41 59.4	+30 32 08	51.3	39.9	14.42	0.02	0.63	0.03	-0.07	0.02	1	2	0.03	0.02

Table 2—Continued

Star	RA	DEC	RA	DEC	l	b	V	σ_V	$B - V$	σ_{B-V}	$U - B$	σ_{U-B}	N	N	$E(B - V)$	S_B
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14) ⁿⁱ	(15) ^{ob}	(16)	(17)
BS 16556−0107	16 45 06.4	+28 57 48	16 47 05.1	+28 52 30	49.5	38.5	13.94	0.01	0.54	0.02	−0.08	0.03	2	6	0.04	0.03
BS 16920−0017	12 04 44.9	+41 56 17	12 07 17.0	+41 39 35	152.0	72.9	13.93	0.04	0.76	0.03	0.15	0.01	1	2	0.01	0.00
BS 16928−0053	12 19 58.6	+34 28 02	12 22 28.1	+34 11 24	162.6	80.6	13.54	0.03	0.84	0.05	0.25	0.02	1	4	0.01	0.00
BS 17436−0058	13 34 39.8	+45 18 23	13 36 46.2	+45 03 07	99.1	69.9	13.83	0.02	0.74	0.01	0.22	0.08	1	2	0.02	0.00
BS 17447−0029	17 48 26.6	+60 12 42	17 49 03.8	+60 11 52	89.0	31.0	13.74	0.01	0.38	0.01	−0.30	0.01	1	6	0.04	0.04
BS 17447−0058	17 58 44.0	+57 21 19	17 59 33.5	+57 21 15	85.8	29.5	14.58	0.04	0.48	0.06	−0.22	0.04	2	4	0.04	0.04
BS 17447−0059	17 58 40.5	+57 21 22	17 59 30.0	+57 21 18	85.8	29.5	14.27	0.01	0.50	0.01	−0.04	0.02	1	2	0.04	0.04
BS 17569−0011	21 55 49.5	+02 39 53	21 58 21.5	+02 54 13	61.8	−38.6	10.75	0.01	1.47	0.01	0.88	0.01	2	8	0.05	0.04
BS 17569−0034	21 59 38.2	+04 18 42	22 02 09.3	+04 33 11	64.1	−38.3	13.28	0.01	0.49	0.01	−0.08	0.01	1	2	0.04	0.04
BS 17569−0037	21 59 00.2	+05 35 28	22 01 30.6	+05 49 55	65.2	−37.3	14.44	0.01	0.44	0.02	−0.12	0.02	2	5	0.05	0.07
BS 17569−0058	22 04 16.1	+06 02 38	22 06 46.3	+06 17 17	66.8	−38.0	12.56	0.01	0.45	0.01	0.00	0.01	1	2	0.08	0.08
BS 17569−0081	22 08 30.3	+06 15 28	22 11 00.5	+06 30 15	67.9	−38.7	13.13	0.04	0.89	0.04	0.45	0.02	1	2	0.09	0.08
BS 17569−0089	22 11 33.5	+03 38 45	22 14 05.2	+03 53 39	66.0	−41.0	13.15	0.01	0.57	0.03	0.00	0.03	1	2	0.06	0.05
BS 17569−0090	22 10 54.4	+04 21 05	22 13 25.7	+04 35 57	66.6	−40.4	14.28	0.03	0.44	0.02	−0.03	0.07	1	2	0.08	0.06
BS 17570−0008	00 04 58.4	+25 52 25	00 07 33.0	+26 09 06	110.8	−35.7	13.70	0.03	0.44	0.04	−0.11	0.03	2	4	0.03	0.02
BS 17570−0009	00 04 33.4	+25 46 11	00 07 07.9	+26 02 52	110.7	−35.8	13.11	0.01	0.53	0.01	−0.01	0.03	2	2	0.03	0.02
BS 17570−0011	00 05 33.5	+25 14 53	00 08 08.1	+25 31 34	110.8	−36.3	12.87	0.01	0.39	0.01	−0.03	0.01	2	3	0.04	0.02
BS 17570−0031	00 11 15.9	+23 34 06	00 13 51.2	+23 50 46	112.0	−38.2	12.30	0.01	0.41	0.02	−0.22	0.01	2	3	0.05	0.03
BS 17570−0041	00 13 28.7	+24 48 17	00 16 04.4	+25 04 56	112.9	−37.1	14.06	0.01	0.40	0.01	0.00	0.01	1	2	0.03	0.02
BS 17570−0042	00 13 50.3	+24 15 48	00 16 26.0	+24 32 27	112.9	−37.6	13.62	0.01	0.45	0.02	−0.19	0.01	2	3	0.04	0.01
BS 17570−0050	00 16 05.9	+26 09 43	00 18 42.1	+26 26 21	113.9	−35.9	13.62	0.01	0.65	0.02	0.15	0.04	1	2	0.04	0.03
BS 17570−0052	00 16 32.7	+26 32 56	00 19 09.0	+26 49 34	114.1	−35.5	12.31	0.02	0.35	0.01	0.06	0.01	1	4	0.03	0.03
BS 17570−0057	00 17 42.7	+26 55 35	00 20 19.2	+27 12 13	114.5	−35.2	13.40	0.01	0.43	0.01	−0.11	0.01	2	4	0.03	0.01
BS 17570−0063	00 17 59.9	+23 30 59	00 20 36.1	+23 47 37	113.9	−38.5	14.51	0.03	0.33	0.01	−0.24	0.02	3	3	0.04	0.02
BS 17570−0067	00 18 45.7	+23 38 59	00 21 22.0	+23 55 36	114.2	−38.4	13.10	0.01	0.37	0.01	0.19	0.01	2	4	0.04	0.02
BS 17570−0071	00 19 57.1	+26 03 38	00 22 33.8	+26 20 15	114.9	−36.1	13.72	0.01	0.44	0.01	−0.02	0.01	1	2	0.03	0.02
BS 17570−0090	00 23 07.1	+22 49 03	00 25 43.8	+23 05 38	115.3	−39.4	13.15	0.02	0.39	0.02	−0.18	0.01	4	6	0.05	0.01
BS 17570−0092	00 23 37.2	+23 42 40	00 26 14.1	+23 59 15	115.6	−38.5	14.01	0.02	0.51	0.02	−0.10	0.03	2	5	0.03	0.02
BS 17570−0093	00 23 20.8	+25 05 35	00 25 57.9	+25 22 10	115.7	−37.1	13.81	0.02	0.44	0.01	−0.11	0.01	2	6	0.03	0.01
BS 17570−0096	00 25 21.2	+25 59 05	00 27 58.7	+26 15 39	116.4	−36.3	12.95	0.01	0.40	0.01	−0.03	0.01	1	2	0.04	0.01
BS 17574−0129	04 47 50.8	+08 55 47	04 50 34.5	+09 00 51	189.7	−21.8	10.72	0.01	0.59	0.01	0.11	0.01	3	4	0.16	0.16
BS 17575−0052	04 53 31.3	+08 49 22	04 56 15.0	+08 54 02	190.7	−20.7	11.31	0.02	0.38	0.01	0.17	0.05	3	3	0.13	0.13
BS 17575−0059	04 54 55.4	+09 00 34	04 57 39.3	+09 05 08	190.7	−20.3	10.58	0.01	0.35	0.01	0.05	0.01	3	3	0.13	0.13
BS 17575−0115	04 59 57.4	+10 14 14	05 02 42.8	+10 18 27	190.3	−18.6	12.19	0.01	0.45	0.01	−0.03	0.01	2	2	0.15	0.18
BS 17575−0128	05 00 06.5	+12 47 25	05 02 54.9	+12 51 37	188.1	−17.1	13.49	0.03	0.53	0.02	0.28	0.02	2	3	0.26	0.31
BS 17575−0129	05 00 35.0	+12 59 24	05 03 23.6	+13 03 34	188.0	−16.9	11.24	0.01	0.40	0.01	0.33	0.01	3	3	0.26	0.30
BS 17575−0142	05 02 15.9	+10 48 03	05 05 02.0	+10 52 06	190.2	−17.8	13.72	0.01	0.46	0.01	0.17	0.01	2	2	0.18	0.21
BS 17575−0143	05 01 43.7	+10 45 02	05 04 29.7	+10 49 08	190.1	−17.9	11.62	0.01	0.45	0.01	0.14	0.01	2	2	0.18	0.20
BS 17575−0144	05 00 38.3	+10 43 50	05 03 24.3	+10 48 00	190.0	−18.2	12.04	0.01	0.36	0.01	0.16	0.02	2	2	0.17	0.20
BS 17575−0163	05 02 59.0	+10 05 14	05 05 44.3	+10 09 14	190.9	−18.0	12.61	0.01	0.44	0.01	0.22	0.01	3	4	0.18	0.20
BS 17575−0168	05 02 59.2	+10 43 54	05 05 45.2	+10 47 54	190.3	−17.7	13.39	0.03	0.55	0.02	−0.14	0.01	4	4	0.17	0.21
BS 17575−0169	05 03 30.5	+10 54 29	05 06 16.7	+10 58 27	190.3	−17.5	12.37	0.01	0.42	0.01	0.21	0.01	3	4	0.17	0.21
BS 17575−0213	05 06 54.4	+11 35 33	05 09 41.5	+11 39 17	190.1	−16.4	12.78	0.02	0.58	0.01	0.09	0.01	1	2	0.38	0.33
BS 17578−0001	21 31 39.5	+17 31 38	21 34 00.5	+17 45 00	70.4	−24.4	10.74	0.01	1.20	0.01	1.05	0.01	1	2	0.11	0.10
BS 17578−0002	21 30 18.6	+17 11 57	21 32 39.7	+17 25 15	69.9	−24.4	11.79	0.01	0.69	0.02	0.25	0.01	3	6	0.10	0.09
BS 17578−0014	21 30 27.0	+13 47 17	21 32 50.7	+14 00 35	67.1	−26.7	12.30	0.01	0.55	0.02	−0.02	0.01	2	4	0.10	0.07
BS 17578−0019	21 32 21.8	+14 55 24	21 34 44.8	+15 08 47	68.4	−26.3	12.51	0.01	0.59	0.01	0.03	0.01	1	4	0.08	0.09
BS 17578−0025	21 33 31.5	+17 29 31	21 35 52.6	+17 42 57	70.7	−24.8	11.56	0.01	0.51	0.01	0.05	0.01	3	6	0.13	0.11
BS 17578−0033	21 35 50.9	+13 31 51	21 38 15.1	+13 45 23	67.9	−27.8	12.19	0.01	0.70	0.01	0.15	0.01	3	6	0.13	0.08
BS 17578−0036	21 36 44.3	+14 11 04	21 39 08.1	+14 24 39	68.6	−27.5	12.36	0.01	0.56	0.01	0.06	0.01	2	6	0.10	0.07
BS 17578−0046	21 37 46.8	+16 51 40	21 40 08.7	+17 05 17	71.0	−25.9	12.33	0.02	0.48	0.02	0.09	0.01	1	2	0.13	0.13
BS 17578−0051	21 38 26.8	+16 17 07	21 40 49.2	+16 30 46	70.6	−26.4	11.93	0.01	0.55	0.01	0.10	0.01	1	2	0.11	0.12
BS 17578−0059	21 41 45.3	+12 48 17	21 44 10.4	+13 02 04	68.3	−29.4	12.37	0.01	0.51	0.01	−0.01	0.01	2	4	0.12	0.09
BS 17578−0060	21 41 02.6	+13 40 31	21 43 27.0	+13 54 17	68.9	−28.7	12.14	0.03	0.52	0.01	−0.17	0.02	4	8	0.09	0.07
BS 17578−0061	21 42 12.7	+13 37 43	21 44 37.2	+13 51 31	69.1	−28.9	11.82	0.02	0.51	0.02	0.03	0.05	3	6	0.09	0.07
BS 17578−0062	21 40 29.5	+13 52 51	21 42 53.7	+14 06 35	69.0	−28.4	11.53	0.04	0.51	0.03	0.25	0.06	2	4	0.12	0.07

Table 2—Continued

Star	RA	DEC	RA	DEC	l	b	V	σ_V	$B - V$	σ_{B-V}	$U - B$	σ_{U-B}	N	N	$E(B - V)$	S_B
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14) ⁿⁱ	(15) ^{ob}	(16)	(17)
BS 17578−0065	21 40 36.2	+15 40 02	21 42 59.2	+15 53 46	70.5	−27.2	12.21	0.01	0.59	0.01	−0.09	0.01	1	2	0.11	0.10
BS 17578−0068	21 40 40.5	+17 06 05	21 43 02.5	+17 19 50	71.7	−26.3	12.98	0.01	0.46	0.01	−0.13	0.01	1	2	0.11	0.10
BS 17578−0071	21 42 48.7	+17 35 43	21 45 10.5	+17 49 33	72.5	−26.3	11.88	0.01	0.42	0.01	−0.03	0.01	1	2	0.11	0.11
BS 17578−0087	21 42 31.3	+12 58 19	21 44 56.3	+13 12 08	68.6	−29.4	11.97	0.01	0.37	0.01	0.09	0.01	1	2	0.11	0.08
BS 17578−0088	21 42 51.9	+12 55 25	21 45 17.0	+13 09 15	68.6	−29.5	12.84	0.01	0.53	0.01	−0.03	0.01	2	6	0.11	0.07
BS 17578−0090	21 45 21.8	+12 48 43	21 47 47.1	+13 02 39	69.0	−30.0	12.62	0.01	0.67	0.01	0.01	0.03	1	2	0.11	0.06
BS 17578−0097	21 48 30.3	+17 09 55	21 50 52.8	+17 23 59	73.2	−27.6	12.52	0.01	0.56	0.01	0.06	0.02	1	2	0.10	0.08
BS 17578−0103	21 47 27.4	+13 46 13	21 49 52.1	+14 00 14	70.2	−29.7	12.95	0.01	0.63	0.03	0.20	0.10	2	4	0.13	0.08
BS 17578−0117	21 50 29.6	+16 18 48	21 52 52.8	+16 32 56	72.9	−28.5	12.55	0.01	0.43	0.01	0.07	0.01	1	4	0.07	0.06
BS 17578−0118	21 50 59.7	+16 19 21	21 53 23.0	+16 33 30	73.0	−28.6	12.15	0.01	0.51	0.01	−0.04	0.04	2	6	0.07	0.06
BS 17579−0011	00 45 18.0	+02 50 33	00 47 52.4	+03 06 53	121.2	−59.7	12.18	0.01	0.52	0.01	−0.09	0.01	1	2	0.02	0.01
BS 17579−0016	00 47 43.3	+00 32 19	00 50 17.2	+00 48 37	122.3	−62.1	12.42	0.01	0.43	0.01	−0.08	0.01	2	6	0.02	0.01
BS 17579−0020	00 49 22.0	+02 17 31	00 51 56.3	+02 33 48	123.2	−60.3	12.47	0.01	0.54	0.01	−0.12	0.01	1	2	0.03	0.02
BS 17579−0033	00 55 32.5	+01 24 47	00 58 06.6	+01 40 57	126.4	−61.1	12.51	0.01	0.42	0.01	−0.12	0.01	2	6	0.03	0.01
BS 17579−0040	00 57 36.3	+03 02 12	01 00 10.9	+03 18 20	127.2	−59.5	13.50	0.03	0.59	0.05	−0.08	0.04	1	2	0.02	0.02
BS 17579−0047	01 03 24.3	+01 40 58	01 05 58.6	+01 56 59	130.4	−60.7	12.44	0.01	0.54	0.02	0.01	0.01	1	2	0.02	0.00
BS 17579−0049	00 55 55.2	+01 47 04	00 58 29.4	+02 03 14	126.5	−60.8	12.43	0.01	0.60	0.01	−0.06	0.01	1	2	0.02	0.01
BS 17579−0050	00 55 40.9	+01 41 38	00 58 15.1	+01 57 48	126.4	−60.9	12.58	0.01	0.59	0.01	0.16	0.01	1	2	0.02	0.01
BS 17579−0051	00 55 34.4	+01 44 14	00 58 08.6	+02 00 24	126.4	−60.8	11.82	0.01	0.49	0.02	−0.09	0.01	1	2	0.02	0.01
BS 17583−0011	21 30 55.1	+25 17 32	21 33 09.7	+25 30 52	76.4	−18.9	11.95	0.01	0.53	0.01	0.03	0.02	1	4	0.11	0.09
BS 17583−0012	21 30 25.1	+25 09 18	21 32 39.8	+25 22 36	76.2	−18.9	12.45	0.01	0.54	0.01	0.10	0.02	1	4	0.11	0.09
BS 17583−0013	21 29 46.7	+24 59 03	21 32 01.5	+25 12 19	75.9	−19.0	12.33	0.01	0.62	0.01	0.05	0.01	1	2	0.14	0.09
BS 17583−0014	21 30 12.6	+24 56 53	21 32 27.4	+25 10 11	76.0	−19.1	12.79	0.01	0.60	0.01	0.14	0.01	1	6	0.14	0.09
BS 17583−0015	21 30 25.8	+24 49 18	21 32 40.8	+25 02 36	75.9	−19.2	11.58	0.01	0.50	0.01	0.11	0.01	1	2	0.13	0.09
BS 17583−0028	21 33 20.2	+24 31 42	21 35 35.7	+24 45 08	76.2	−19.9	12.98	0.02	0.55	0.01	0.03	0.04	2	4	0.11	0.09
BS 17583−0032	21 31 45.4	+26 22 43	21 33 59.2	+26 36 05	77.3	−18.3	11.46	0.01	0.54	0.01	0.23	0.02	1	2	0.14	0.11
BS 17583−0057	21 35 55.1	+24 24 17	21 38 11.0	+24 37 49	76.5	−20.4	13.05	0.01	0.58	0.01	−0.02	0.02	1	2	0.07	0.06
BS 17583−0064	21 38 29.7	+27 25 46	21 40 43.4	+27 39 25	79.2	−18.6	12.94	0.01	0.51	0.02	0.16	0.05	2	4	0.11	0.09
BS 17583−0065	21 39 57.6	+27 28 45	21 42 11.4	+27 42 28	79.5	−18.8	12.59	0.01	0.37	0.01	0.08	0.02	2	4	0.10	0.08
BS 17583−0067	21 38 52.5	+27 06 24	21 41 06.5	+27 20 04	79.0	−18.9	12.56	0.02	0.39	0.01	0.29	0.03	2	6	0.11	0.09
BS 17583−0070	21 38 58.9	+26 19 46	21 41 13.6	+26 33 26	78.5	−19.5	12.45	0.01	0.36	0.10	0.15	0.14	1	2	0.10	0.08
BS 17583−0077	21 39 31.3	+24 50 57	21 41 47.2	+25 04 39	77.5	−20.6	12.36	0.01	0.51	0.01	0.03	0.02	2	6	0.07	0.06
BS 17583−0083	21 38 23.7	+23 38 01	21 40 40.5	+23 51 40	76.4	−21.3	12.46	0.01	0.47	0.01	−0.21	0.15	1	2	0.07	0.07
BS 17583−0084	21 39 16.8	+23 10 43	21 41 34.0	+23 24 24	76.2	−21.8	12.86	0.01	0.55	0.01	0.06	0.01	1	2	0.10	0.08
BS 17583−0092	21 40 11.2	+24 26 05	21 42 27.5	+24 39 48	77.3	−21.0	10.02	0.01	0.34	0.01	0.13	0.02	2	6	0.09	0.07
BS 17583−0094	21 40 21.3	+25 38 29	21 42 36.7	+25 52 13	78.2	−20.2	12.54	0.01	0.65	0.01	0.10	0.01	1	2	0.09	0.08
BS 17583−0095	21 41 26.8	+25 48 30	21 43 42.2	+26 02 16	78.5	−20.2	13.25	0.02	0.52	0.01	0.01	0.02	1	4	0.09	0.06
BS 17583−0097	21 42 04.0	+25 59 20	21 44 19.3	+26 13 08	78.8	−20.2	12.13	0.01	0.71	0.01	0.17	0.01	1	3	0.09	0.06
BS 17583−0110	21 42 27.0	+26 30 01	21 44 41.9	+26 43 50	79.2	−19.9	10.99	0.01	0.27	0.01	0.16	0.01	1	4	0.09	0.07
BS 17583−0116	21 43 02.4	+24 34 47	21 45 18.9	+24 48 37	77.9	−21.4	12.22	0.01	0.66	0.01	−0.02	0.01	1	4	0.07	0.07
BS 17583−0124	21 43 44.7	+23 22 29	21 46 02.3	+23 36 21	77.1	−22.4	12.52	0.01	0.50	0.01	0.00	0.01	1	2	0.09	0.07
BS 17583−0125	21 43 17.9	+23 14 01	21 45 35.5	+23 27 52	76.9	−22.4	12.74	0.01	0.63	0.01	0.13	0.01	1	4	0.10	0.07
BS 17583−0128	21 44 36.0	+23 14 58	21 46 53.7	+23 28 52	77.2	−22.6	13.01	0.01	0.49	0.01	0.05	0.01	1	2	0.09	0.07
BS 17583−0132	21 46 31.1	+23 17 38	21 48 49.0	+23 31 37	77.6	−22.9	12.06	0.01	0.63	0.01	0.08	0.01	1	2	0.08	0.07
BS 17583−0134	21 44 39.1	+23 48 32	21 46 56.4	+24 02 26	77.6	−22.2	12.10	0.01	0.41	0.01	−0.02	0.01	1	6	0.07	0.07
BS 17583−0140	21 46 37.8	+26 30 11	21 48 53.2	+26 44 10	79.9	−20.5	12.31	0.02	0.57	0.01	0.00	0.01	1	2	0.07	0.07
BS 17583−0146	21 47 45.6	+26 00 40	21 50 01.6	+26 14 42	79.8	−21.1	12.19	0.01	0.49	0.01	−0.02	0.02	1	2	0.07	0.07
BS 17583−0152	21 47 32.1	+24 45 11	21 49 49.0	+24 59 12	78.8	−21.9	12.20	0.01	0.57	0.01	0.07	0.01	1	4	0.07	0.07
BS 17583−0154	21 47 52.8	+24 17 04	21 50 10.1	+24 31 06	78.6	−22.3	12.72	0.01	0.44	0.01	−0.01	0.01	1	4	0.06	0.05
BS 17583−0155	21 47 44.7	+23 45 23	21 50 02.4	+23 59 25	78.1	−22.7	12.31	0.01	0.49	0.01	0.09	0.02	1	2	0.06	0.06
BS 17583−0157	21 46 47.6	+22 44 29	21 49 06.0	+22 58 28	77.2	−23.3	12.17	0.02	0.81	0.01	0.35	0.01	3	6	0.08	0.07
BS 17583−0160	21 49 06.4	+23 37 38	21 51 24.3	+23 51 43	78.3	−23.0	12.89	0.01	0.61	0.01	0.17	0.01	1	2	0.06	0.06
BS 17583−0163	21 49 43.0	+24 29 50	21 52 00.4	+24 43 56	79.0	−22.5	12.28	0.01	0.51	0.01	0.04	0.01	1	2	0.06	0.05
BS 17583−0165	21 48 57.3	+25 07 48	21 51 14.1	+25 21 52	79.4	−21.9	12.79	0.01	0.46	0.01	−0.01	0.02	1	2	0.07	0.07
BS 17584−0005	23 20 31.6	+24 15 08	23 23 00.3	+24 31 35	98.5	−34.1	11.79	0.01	0.54	0.02	−0.03	0.02	2	4	0.06	0.04
BS 17584−0012	23 21 46.5	+23 55 48	23 24 15.4	+24 12 16	98.7	−34.5	11.31	0.01	1.10	0.01	0.68	0.03	1	2	0.05	0.03
BS 17584−0016	23 21 55.8	+26 09 38	23 24 24.2	+26 26 06	99.8	−32.5	12.06	0.02	0.65	0.02	0.08	0.03	1	2	0.07	0.05

Table 2—Continued

Star	RA	DEC	RA	DEC	l	b	V	σ_V	$B - V$	σ_{B-V}	$U - B$	σ_{U-B}	N	N	$E(B - V)$	
(1)	(2)	1950 (3)	(4)	2000 (5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14) ⁿⁱ	(15) ^{ob}	S _B (16)	(17)
BS 17584−0021	23 23 51.7	+26 41 19	23 26 20.3	+26 57 48	100.5	−32.2	12.74	0.03	0.51	0.01	−0.05	0.08	2	4	0.08	0.04
BS 17584−0030	23 30 00.3	+26 50 11	23 32 29.7	+27 06 44	102.0	−32.5	11.96	0.01	0.40	0.02	0.09	0.02	2	4	0.07	0.04
BS 17584−0032	23 29 40.3	+26 25 14	23 32 09.8	+26 41 47	101.8	−32.9	12.92	0.02	0.57	0.01	−0.11	0.01	2	4	0.07	0.04
BS 17584−0035	23 28 48.7	+24 39 14	23 31 18.4	+24 55 47	100.8	−34.5	11.93	0.01	0.75	0.01	0.01	0.03	1	2	0.09	0.04
BS 17584−0037	23 29 55.5	+23 49 01	23 32 25.5	+24 05 34	100.7	−35.3	12.25	0.01	0.67	0.01	0.29	0.02	1	4	0.06	0.03
BS 17584−0038	23 30 08.1	+23 46 10	23 32 38.1	+24 02 43	100.7	−35.4	12.30	0.01	0.70	0.02	0.07	0.05	2	5	0.06	0.03
BS 17584−0039	23 31 25.0	+22 51 13	23 33 55.3	+23 07 47	100.7	−36.4	12.35	0.01	0.52	0.01	0.02	0.01	1	2	0.08	0.03
BS 17584−0040	23 31 30.7	+23 24 21	23 34 00.9	+23 40 55	100.9	−35.9	12.40	0.01	0.51	0.01	0.05	0.02	1	2	0.07	0.04
BS 17584−0041	23 31 20.5	+24 03 34	23 33 50.6	+24 20 08	101.2	−35.2	12.57	0.01	0.58	0.01	0.03	0.01	1	2	0.08	0.03
BS 17584−0043	23 32 00.1	+24 50 06	23 34 30.2	+25 06 40	101.7	−34.6	12.53	0.01	0.47	0.01	0.08	0.02	1	2	0.06	0.03
BS 17584−0045	23 30 34.1	+25 16 40	23 33 03.9	+25 33 14	101.5	−34.0	12.89	0.01	0.44	0.01	0.03	0.01	1	4	0.04	0.03
BS 17584−0046	23 31 27.7	+25 57 54	23 33 57.5	+26 14 28	102.0	−33.5	12.74	0.01	0.36	0.01	−0.18	0.01	1	2	0.05	0.03
BS 17584−0048	23 30 31.0	+26 24 46	23 33 00.6	+26 41 19	102.0	−33.0	11.18	0.01	0.45	0.01	0.05	0.01	1	2	0.08	0.04
BS 17584−0051	23 33 12.5	+26 53 00	23 35 42.4	+27 09 35	102.9	−32.7	9.64	0.01	0.44	0.01	−0.23	0.01	1	2	0.06	0.04
BS 17584−0053	23 32 50.7	+26 27 55	23 35 20.6	+26 44 30	102.6	−33.1	10.86	0.01	0.34	0.01	−0.10	0.01	1	2	0.06	0.03
BS 17584−0061	23 35 15.5	+25 25 13	23 37 45.9	+25 41 49	102.8	−34.3	12.85	0.01	0.50	0.01	0.01	0.01	1	2	0.06	0.03
BS 17584−0070	23 37 13.6	+25 20 14	23 39 44.3	+25 36 51	103.3	−34.5	12.36	0.03	0.52	0.01	−0.18	0.06	2	4	0.04	0.03
BS 17584−0071	23 38 46.2	+24 01 32	23 41 17.3	+24 18 09	103.1	−35.9	12.82	0.01	0.55	0.01	−0.02	0.01	1	2	0.05	0.02
BS 17584−0074	23 36 45.5	+22 58 25	23 39 16.5	+23 15 01	102.2	−36.7	9.74	0.01	0.35	0.01	−0.10	0.01	1	2	0.04	0.02
BS 17584−0083	23 39 47.0	+25 00 33	23 42 18.1	+25 17 11	103.8	−35.0	12.77	0.01	0.54	0.01	0.01	0.01	1	2	0.04	0.02
BS 17584−0084	23 39 20.6	+25 02 10	23 41 51.6	+25 18 47	103.7	−34.9	13.21	0.03	0.50	0.02	−0.19	0.01	1	2	0.04	0.02
BS 17584−0088	23 33 16.8	+25 59 00	23 35 46.9	+26 15 35	102.5	−33.6	13.40	0.01	0.51	0.01	−0.21	0.03	1	2	0.04	0.02
BS 17585−0132	04 29 21.0	+16 12 35	04 32 12.7	+16 18 55	180.5	−21.1	12.38	0.01	0.69	0.02	0.41	0.02	3	3	0.39	0.27
CS 22179−0024	00 42 07.4	−03 07 43	00 44 40.4	−02 51 19	118.8	−65.7	14.14	0.04	0.39	0.03	−0.25	0.03	2	3	0.03	0.04
CS 22872−0036	16 20 08.6	−02 16 59	16 22 44.8	−02 23 56	11.5	31.2	14.88	0.03	0.51	0.04	−0.11	0.04	2	3	0.15	0.13
CS 22889−0045	13 41 07.7	−08 25 23	13 43 45.7	−08 40 26	324.3	52.0	13.31	0.05	0.46	0.02	−0.32	0.01	1	2	0.04	0.02
CS 22962−0021	01 46 34.7	−04 05 46	01 49 06.3	−03 50 53	156.0	−63.0	13.46	0.01	0.50	0.02	−0.19	0.02	1	1	0.02	0.02
CS 22962−0027	01 45 44.1	−06 09 18	01 48 14.6	−05 54 23	158.0	−64.8	14.56	0.01	0.52	0.04	−0.02	0.06	1	2	0.03	0.00
CS 22965−0016	21 54 31.2	−04 48 25	21 57 07.7	−04 34 06	53.5	−42.8	14.07	0.02	0.74	0.02	0.05	0.04	2	5	0.04	0.03
CS 22965−0029	21 56 56.1	−03 51 00	21 59 32.0	−03 36 36	55.1	−42.7	13.80	0.01	0.60	0.04	−0.02	0.04	2	4	0.08	0.06
CS 22965−0053	21 59 50.0	−02 38 04	22 02 25.1	−02 23 34	57.0	−42.6	12.75	0.01	0.43	0.02	0.19	0.03	1	2	0.13	0.06
CS 22965−0054	22 03 54.8	−02 47 18	22 06 30.0	−02 32 39	57.6	−43.5	15.10	0.06	0.59	0.02	−0.10	0.11	1	2	0.13	0.15
CS 29502−0001	22 27 20.0	−02 02 16	22 29 54.6	−01 46 52	63.7	−47.7	11.40	0.01	0.42	0.01	−0.01	0.01	1	2	0.06	0.04
CS 29502−0009	22 24 50.2	+01 43 16	22 27 23.0	+01 58 34	67.1	−44.7	13.83	0.01	0.71	0.03	−0.06	0.05	1	3	0.08	0.04
CS 29502−0042	22 19 16.1	+02 13 39	22 21 48.6	+02 28 47	66.3	−43.4	12.78	0.02	0.60	0.02	−0.03	0.02	1	2	0.07	0.04
CS 29502−0062	22 12 49.6	+02 16 09	22 15 22.0	+02 31 05	64.9	−42.1	11.30	0.01	0.50	0.01	−0.02	0.02	1	2	0.05	0.03
CS 29502−0067	22 12 05.2	−01 44 11	22 14 39.7	−01 29 15	60.5	−44.5	13.94	0.01	0.77	0.04	−0.02	0.04	1	2	0.11	0.05
CS 29516−0017	22 25 26.8	+03 44 08	22 27 58.7	+03 59 27	69.3	−43.5	12.41	0.01	0.08	0.01	0.13	0.01	1	2	0.10	0.06
CS 29516−0024	22 23 42.7	+02 36 33	22 26 15.1	+02 51 49	67.8	−43.9	13.57	0.01	0.86	0.01	0.20	0.01	1	2	0.10	0.06
CS 29516−0026	22 24 23.1	+02 48 08	22 26 55.4	+03 03 25	68.1	−43.9	12.93	0.01	0.34	0.02	0.23	0.01	1	2	0.11	0.06
CS 29516−0028	22 23 09.2	+05 22 24	22 25 40.3	+05 37 39	70.4	−41.9	15.02	0.03	0.55	0.07	−0.28	0.02	1	2	0.13	0.06
CS 29516−0032	22 21 47.7	+06 02 47	22 24 18.4	+06 18 00	70.7	−41.2	14.53	0.01	0.66	0.04	0.08	0.05	1	2	0.10	0.06
CS 29516−0037	22 22 55.8	+04 17 57	22 25 27.4	+04 33 12	69.3	−42.6	12.32	0.01	0.23	0.01	0.25	0.01	1	2	0.14	0.06
CS 29516−0096	22 11 57.5	+05 23 37	22 14 28.3	+05 38 31	67.8	−39.9	15.31	0.03	0.17	0.01	0.16	0.07	1	2	0.10	0.07
CS 29521−0012	22 50 29.6	+09 00 22	22 53 00.1	+09 16 19	80.6	−43.6	14.82	0.02	0.60	0.01	−0.20	0.07	1	2	0.06	0.04
CS 29521−0031	22 58 54.6	+09 17 46	23 01 25.4	+09 33 53	83.1	−44.6	12.81	0.01	0.61	0.01	−0.04	0.01	1	4	0.07	0.03
CS 29521−0036	23 00 20.7	+08 43 15	23 02 51.8	+08 59 24	83.1	−45.3	12.54	0.02	0.47	0.02	−0.04	0.01	1	2	0.07	0.03
CS 29521−0065	23 00 39.5	+10 31 21	23 03 10.0	+10 47 30	84.6	−43.9	11.24	0.01	0.30	0.01	0.20	0.01	1	2	0.06	0.03
CS 29522−0046	23 42 26.6	+08 30 14	23 44 59.6	+08 46 53	96.5	−50.6	12.74	0.01	0.49	0.03	−0.13	0.03	3	5	0.11	0.06
CS 30312−0006	15 30 22.2	−01 43 13	15 32 57.6	−01 53 16	2.8	41.5	13.64	0.01	0.62	0.01	−0.05	0.01	2	3	0.14	0.08
CS 30312−0044	15 34 34.3	+00 42 42	15 37 07.4	+00 32 52	6.2	42.2	13.34	0.01	0.78	0.03	0.09	0.02	2	3	0.08	0.06
CS 30312−0067	15 33 00.3	−02 02 17	15 35 36.0	−02 12 11	3.0	40.8	14.23	0.02	0.57	0.03	0.11	0.04	2	5	0.13	0.10
CS 30325−0023	15 02 50.8	+04 19 07	15 05 20.9	+04 07 31	3.1	50.6	13.93	0.03	0.73	0.03	0.12	0.04	1	2	0.04	0.02
CS 30325−0033	15 02 59.8	+07 12 05	15 05 27.5	+07 00 29	6.8	52.4	12.94	0.01	0.44	0.02	−0.21	0.01	1	3	0.03	0.01
CS 30332−0118	22 31 50.7	+10 00 05	22 34 20.0	+10 15 35	76.6	−40.0	14.35	0.02	0.46	0.03	−0.16	0.02	1	3	0.10	0.06
CS 30332−0119	22 32 38.7	+10 10 29	22 35 07.9	+10 26 00	77.0	−40.0	15.28	0.04	0.63	0.06	0.12	0.05	1	3	0.09	0.06

Table 2—Continued

Star	RA	DEC	RA	DEC	l	b	V	σ_V	$B - V$	σ_{B-V}	$U - B$	σ_{U-B}	N	N	$E(B - V)$	
(1)	(2)	1950 (3)	(4)	2000 (5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14) ⁿⁱ	(15) ^{ob}	S (16)	B (17)
CS 30332-0138	22 30 44.3	+09 53 44	22 33 13.6	+10 09 12	76.3	-39.9	13.98	0.03	1.19	0.02	0.83	0.08	1	2	0.10	0.06
CS 30333-0002	23 47 27.6	+08 26 49	23 50 00.8	+08 43 29	98.3	-51.2	11.32	0.01	0.50	0.01	-0.31	0.01	3	5	0.10	0.05
CS 31061-0004	02 43 34.8	+03 25 45	02 46 11.2	+03 38 18	169.4	-48.6	13.55	0.01	0.60	0.03	0.00	0.06	1	2	0.04	0.03
CS 31061-0032	02 36 07.0	+03 06 08	02 38 43.1	+03 19 02	167.5	-50.0	13.90	0.01	0.40	0.01	-0.28	0.05	2	2	0.04	0.02
CS 31061-0033	02 33 54.9	+02 22 46	02 36 30.4	+02 35 46	167.5	-51.0	13.64	0.01	0.68	0.01	0.00	0.02	1	2	0.03	0.01
CS 31061-0057	02 33 20.2	+02 33 26	02 35 55.8	+02 46 28	167.1	-50.9	14.21	0.03	0.34	0.04	-0.04	0.03	2	3	0.03	0.01
CS 31061-0062	02 30 12.3	+05 31 52	02 32 50.0	+05 45 02	163.4	-49.0	13.90	0.03	0.48	0.03	-0.30	0.04	2	3	0.06	0.04
CS 31061-0072	02 26 37.9	+04 05 53	02 29 14.6	+04 19 13	163.6	-50.7	13.99	0.02	0.48	0.02	-0.27	0.02	2	5	0.04	0.02
CS 31061-0074	02 26 44.3	+04 24 58	02 29 21.2	+04 38 17	163.3	-50.4	13.98	0.01	0.45	0.01	-0.23	0.01	1	2	0.05	0.03
CS 31067-0019	03 20 45.2	+04 28 16	03 23 23.0	+04 38 52	177.9	-41.5	12.61	0.02	0.45	0.01	-0.26	0.01	3	5	0.11	0.09
CS 31067-0039	03 13 03.8	+05 45 42	03 15 42.6	+05 56 44	174.9	-41.9	13.69	0.03	0.87	0.03	0.35	0.05	2	4	0.20	0.14
CS 31067-0045	03 13 26.6	+03 15 17	03 16 03.2	+03 26 18	177.4	-43.6	13.87	0.01	0.69	0.01	0.03	0.05	1	2	0.12	0.07
CS 31069-0005	00 23 10.4	+03 58 50	00 25 44.6	+04 15 25	110.8	-58.0	13.52	0.01	0.57	0.01	0.07	0.01	1	2	0.02	0.00
CS 31069-0082	00 11 47.0	+06 49 56	00 14 21.2	+07 06 35	106.9	-54.6	14.06	0.02	0.47	0.01	-0.06	0.01	1	2	0.06	0.04
CS 31069-0109	00 08 09.4	+06 10 30	00 10 43.4	+06 27 10	105.1	-55.0	12.90	0.01	0.87	0.01	0.50	0.01	1	2	0.05	0.04
CS 31070-0047	00 13 01.8	+02 55 23	00 15 35.7	+03 12 02	105.6	-58.5	14.11	0.05	0.38	0.02	-0.02	0.01	1	2	0.02	0.02
CS 31074-0005	02 52 15.1	+08 39 35	02 54 55.9	+08 51 42	167.0	-43.2	14.21	0.05	0.67	0.05	0.00	0.05	1	1	0.18	0.13
CS 31079-0004	03 02 28.2	+03 40 46	03 05 05.1	+03 52 22	174.3	-45.3	13.15	0.01	0.46	0.03	-0.17	0.01	2	2	0.09	0.08
CS 31084-0022	03 27 17.4	+09 15 58	03 29 59.8	+09 26 12	174.9	-37.0	11.56	0.01	0.45	0.01	-0.17	0.01	2	2	0.22	0.20

Table 3. Photometric Data for Suspected Variable Stars

Star	RA	DEC	RA	DEC	l	b	V	σ_V	$B - V$	σ_{B-V}	$U - B$	σ_{U-B}	N	N	$E(B - V)$	
	1950		2000										N	N	$E(B - V)$	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
BS 16083-0172	14 48 59.5	+48 57 33	14 50 41.6	+48 45 14	84.2	58.4	13.13	0.39	0.61	0.11	0.09	0.01	2	4	0.03	0.01
BS 16542-0002	16 18 24.8	+22 50 18	16 20 33.1	+22 43 12	39.7	42.7	12.62	0.08	0.76	0.04	-0.04	0.04	6	15	0.06	0.04
BS 16556-0081	16 42 37.8	+31 32 02	16 44 32.8	+31 26 34	52.6	39.6	14.32	0.09	0.48	0.08	-0.25	0.02	3	6	0.03	0.02
BS 17569-0014	21 55 34.8	+06 20 16	21 58 04.6	+06 34 36	65.3	-36.2	12.99	0.11	0.48	0.10	-0.11	0.05	1	2	0.06	0.05
BS 17569-0023	21 57 07.8	+05 54 18	21 59 37.9	+06 08 41	65.2	-36.8	11.05	0.38	1.44	0.03	1.39	0.04	2	4	0.05	0.04
BS 17569-0027	21 56 59.5	+03 38 20	21 59 31.0	+03 52 43	63.0	-38.2	13.98	0.13	0.49	0.06	-0.10	0.04	4	11	0.05	0.04
BS 17569-0028	21 58 10.1	+03 26 17	22 00 41.7	+03 40 43	63.0	-38.6	13.89	0.18	0.56	0.06	-0.17	0.03	4	8	0.05	0.03
BS 17569-0047	22 01 06.1	+04 39 20	22 03 37.1	+04 53 52	64.8	-38.4	14.37	0.10	0.47	0.04	-0.16	0.04	2	4	0.05	0.04
BS 17569-0063	22 04 44.6	+07 07 48	22 07 14.2	+07 22 28	67.9	-37.4	15.07	0.28	-0.34	0.18	-1.26	0.07	2	4	0.08	0.07
BS 17569-0078	22 07 54.2	+03 49 25	22 10 25.8	+04 04 11	65.4	-40.2	13.88	0.26	0.51	0.04	-0.16	0.04	3	6	0.04	0.04
BS 17575-0182	05 06 02.6	+12 10 41	05 08 50.4	+12 14 28	189.5	-16.2	12.85	0.09	0.73	0.02	0.43	0.37	3	3	0.38	0.37
BS 17578-0013	21 32 08.1	+13 51 28	21 34 31.9	+14 04 51	67.5	-26.9	12.27	0.15	0.56	0.02	-0.15	0.07	2	4	0.09	0.07
BS 17578-0032	21 34 32.2	+13 37 53	21 36 56.3	+13 51 22	67.7	-27.5	12.12	0.11	0.47	0.01	-0.25	0.07	2	5	0.11	0.07
BS 17578-0056	21 38 44.3	+13 58 07	21 41 08.4	+14 11 47	68.8	-28.1	9.92	0.12	0.38	0.03	0.14	0.06	9	24	0.11	0.06
BS 17578-0106	21 49 47.8	+13 18 52	21 52 13.0	+13 32 59	70.3	-30.5	12.13	0.38	0.58	0.07	0.11	0.07	3	6	0.10	0.07
BS 17579-0012	00 45 24.7	+03 21 24	00 47 59.2	+03 37 44	121.3	-59.2	12.28	0.09	0.10	0.03	-0.91	0.02	2	4	0.02	0.02
BS 17583-0100	21 40 13.1	+26 26 50	21 42 27.8	+26 40 33	78.8	-19.6	12.37	0.10	0.51	0.04	-0.21	0.08	7	20	0.10	0.08
BS 17583-0111	21 43 09.9	+26 16 44	21 45 25.1	+26 30 35	79.2	-20.2	12.63	0.13	0.49	0.04	-0.08	0.07	4	13	0.09	0.07
BS 17583-0115	21 44 00.3	+24 56 11	21 46 16.7	+25 10 04	78.3	-21.3	12.80	0.17	0.67	0.05	0.11	0.06	4	8	0.08	0.06
BS 17584-0024	23 25 16.8	+24 36 28	23 27 46.0	+24 52 58	99.9	-34.2	12.79	0.11	0.56	0.03	0.14	0.04	4	13	0.11	0.04
BS 17584-0034	23 29 40.3	+26 25 14	23 32 09.8	+26 41 47	101.3	-34.2	13.30	0.16	0.49	0.02	-0.03	0.03	2	6	0.05	0.03
CS 22954-0004	02 35 45.6	-05 56 15	02 38 15.0	-05 43 19	177.6	-56.8	14.27	0.15	0.42	0.12	-0.22	0.01	3	4	0.03	0.01
CS 30332-0012	22 48 15.7	+10 19 59	22 50 45.7	+10 35 53	81.1	-42.3	14.65	0.11	0.54	0.05	-0.21	0.04	1	3	0.09	0.04

Table 4. Program Stars With Duplicate HK Survey Identifications

Cataloged Star Name	Alternate Name 1	Alternate Name 2
BS 16034–0115	BS 16079–0069	BS 16084–0001
BS 16076–0012	BS 16942–0037	...
BS 16077–0077	BS 16081–0010	...
BS 16079–0133	BS 16084–0075	...
BS 16082–0129	BS 16078–0085	...
BS 16083–0043	BS 15620–0020	...
CS 29502–0042	CS 29516–0041	...
CS 31061–0032	CS 31063–0035	...
CS 31061–0062	CS 31063–0069	...
CS 31061–0072	CS 31063–0092	...
CS 31079–0004	CS 31078–0002	...
CS 31084–0022	CS 31083–0022	...

Table 5. De-reddened Colors, Abundances, and Classifications

Star (1)	$E(B - V)_A$ (2)	V_o (3)	$(B - V)_o$ (4)	$(U - B)_o$ (5)	$[\text{Fe}/\text{H}]_{\text{K3}}$ (6)	σ_{Fe} (7)	Class (8)	Comment (9)
BS 15622–0053	0.01	13.56	0.73	0.37	...	...	...	CORE EM
BS 15622–0055	0.01	14.02	0.39	−0.18	−1.96:	0.18	TO	
BS 15624–0040	0.01	13.54	0.40	−0.23	−1.90	0.15	TO	
BS 16034–0101	0.01	12.33	0.42	−0.05	−2.12	0.15	FHB	
BS 16034–0115	0.02	11.93	0.35	−0.27	−1.77	0.22	TO	
BS 16076–0006	0.03	13.27	0.49	−0.21	−2.84	0.30	SG	
BS 16076–0012	0.02	13.70	0.45	−0.27	−2.02	0.15	TO	
BS 16076–0040	0.04	11.88	0.36	−0.30	−1.77	0.22	TO	
BS 16076–0050	0.04	12.99	0.52	−0.28	−2.31	0.15	SG	
BS 16077–0007	0.02	12.15	0.31	−0.29	−2.45	0.23	TO	
BS 16077–0023	0.02	12.18	0.35	−0.28	−2.50	0.20	TO	
BS 16077–0047	0.02	14.34	0.51	−0.31	−2.40	0.17	SG	
BS 16077–0077	0.03	13.61	0.58	−0.16	−2.49	0.20	G	
BS 16078–0025	0.02	12.55	0.40	−0.22	−1.76	0.15	TO	BLUE
BS 16079–0019	0.02	14.03	0.36	−0.24	−2.32	0.22	TO	
BS 16079–0133	0.01	12.03	0.42	−0.01	−1.48	0.15	FHB	
BS 16080–0007	0.01	13.43	0.45	−0.23	−1.54	0.15	TO	
BS 16080–0022	0.02	13.07	0.41	−0.23	−1.67	0.15	TO	
BS 16080–0054	0.03	12.70	0.68	0.18	−2.62	0.15	G	
BS 16080–0067	0.01	13.12	0.53	−0.07	−0.67	0.21	TO	
BS 16080–0093	0.02	13.55	0.67	0.12	−2.82	0.22	G	
BS 16080–0175	0.03	14.26	0.42	−0.18	−1.84	0.15	TO	
BS 16082–0029	0.01	13.14	0.58	0.05	−2.45	0.20	G	
BS 16082–0045	0.02	13.25	0.38	−0.27	−1.54	0.15	TO	
BS 16082–0097	0.02	12.06	0.34	−0.19	−1.29	0.22	TO	
BS 16082–0129	0.02	13.55	0.60	0.03	−2.40:	0.15	G	BLUE
BS 16083–0020	0.02	13.58	0.29	−0.22	−2.22::	0.26	FHB/A	
BS 16083–0024	0.04	14.22	0.44	−0.10	−2.75	0.24	FHB	
BS 16083–0043	0.02	13.10	0.37	−0.30	−2.50	0.20	TO	
BS 16083–0065	0.02	14.74	0.42	−0.19	−2.69	0.20	TO	
BS 16083–0146	0.03	14.71	0.25	−0.31	−2.22::	0.26	FHB/A	
BS 16084–0123	0.01	13.53	0.41	−0.02	−0.38	0.17	TO	
BS 16084–0160	0.01	13.09	0.81	0.27	−3.22	0.15	G	
BS 16089–0013	0.01	13.30	0.69	0.07	−2.59	0.15	G	
BS 16089–0042	0.02	14.14	0.38	−0.02	−1.87	0.17	FHB	
BS 16089–0086	0.02	14.05	0.30	0.09	−1.68	0.26	FHB	
BS 16468–0029	0.02	13.48	0.39	−0.28	−0.57	0.16	TO	
BS 16542–0016	0.06	13.38	0.67	−0.36	−0.46	0.35	SG	CORE EM
BS 16542–0022	0.05	12.80	0.38	−0.31	−2.29	0.18	TO	
BS 16542–0051	0.05	14.37	0.47	−0.08	−2.24	0.15	FHB	
BS 16542–0052	0.05	12.61	0.82	0.38	...	...	...	
BS 16545–0074	0.02	14.03	0.41	−0.28	−2.54	0.17	TO	
BS 16545–0082	0.02	12.96	0.47	−0.22	−2.37	0.15	SG	
BS 16545–0089	0.02	14.32	0.36	−0.31	−3.42	0.17	TO	
BS 16548–0028	0.03	14.56	0.40	−0.21	−1.76	0.15	TO	
BS 16548–0070	0.02	13.63	0.65	−0.07	−2.35	0.15	G	
BS 16548–0140	0.03	12.53	0.44	−0.21	−2.10	0.15	TO	
BS 16549–0017	0.02	12.36	0.37	−0.26	−2.32	0.22	TO	
BS 16552–0038	0.02	14.54	0.52	−0.13	−2.26	0.15	SG	
BS 16552–0042	0.02	14.58	0.37	0.03	−2.50	0.20	FHB	
BS 16552–0086	0.02	14.54	0.60	0.14	...	...	...	BLUE
BS 16552–0090	0.02	13.22	0.72	0.17	−1.11	0.37	G	
BS 16552–0142	0.02	11.98	0.44	−0.28	−2.35	0.15	TO	
BS 16554–0053	0.03	14.96	0.26	−0.28	−1.38::	0.24	FHB/A	
BS 16556–0045	0.03	13.85	0.61	0.00	−0.50	0.31	TO	
BS 16556–0046	0.02	14.12	0.54	−0.11	−1.01	0.20	TO	
BS 16556–0054	0.03	14.33	0.60	−0.09	−1.84	0.22	SG	
BS 16556–0107	0.04	13.82	0.50	−0.11	−1.44	0.15	TO	
BS 16920–0017	0.01	13.90	0.75	0.14	−3.28	0.16	G	
BS 16928–0053	0.01	13.51	0.83	0.24	−3.10	0.15	G	

Table 5—Continued

Star (1)	$E(B - V)_A$ (2)	V_o (3)	$(B - V)_o$ (4)	$(U - B)_o$ (5)	$[\text{Fe}/\text{H}]_{\text{K3}}$ (6)	σ_{Fe} (7)	Class (8)	Comment (9)
BS 17436–0058	0.02	13.77	0.72	0.21	–2.41	0.15	G	
BS 17447–0029	0.04	13.62	0.34	–0.33	–2.79	0.19	TO	
BS 17447–0058	0.04	14.46	0.44	–0.25	–0.71	0.15	TO	
BS 17447–0059	0.04	14.15	0.46	–0.07	–0.77	0.15	TO	
BS 17569–0011	0.05	10.60	1.42	0.85	...	...	...	CORE EM
BS 17569–0034	0.04	13.16	0.45	–0.11	–1.45:	0.15	TO/FHB	
BS 17569–0037	0.05	14.29	0.39	–0.16	–1.54	0.15	TO	
BS 17569–0058	0.07	12.34	0.38	–0.05	–1.34:	0.15	TO/FHB	
BS 17569–0081	0.09	12.85	0.80	0.39	...	...	...	CORE EM
BS 17569–0089	0.06	12.97	0.51	–0.04	–0.34	0.18	TO	
BS 17569–0090	0.08	14.03	0.36	–0.09	+0.01	0.25	TO	
BS 17570–0008	0.03	13.61	0.41	–0.13	–0.26	0.18	TO	
BS 17570–0009	0.03	13.02	0.50	–0.03	–2.06	0.15	SG	
BS 17570–0011	0.04	12.75	0.35	–0.06	–2.32	0.22	FHB	
BS 17570–0031	0.04	12.17	0.37	–0.25	–1.66	0.21	TO	
BS 17570–0041	0.03	13.97	0.37	–0.02	–0.74:	0.17	TO/FHB	
BS 17570–0042	0.04	13.50	0.41	–0.22	–1.67	0.15	TO	
BS 17570–0050	0.04	13.50	0.61	0.12	–1.18	0.30	SG	
BS 17570–0052	0.03	12.22	0.32	0.04	–0.81	0.18	FHB	
BS 17570–0057	0.03	13.31	0.40	–0.13	–1.58:	0.15	TO/FHB	
BS 17570–0063	0.04	14.39	0.29	–0.27	–2.91::	0.19	FHB/A	BLUE
BS 17570–0067	0.04	12.98	0.33	0.16	–1.29	0.22	FHB	
BS 17570–0071	0.03	13.63	0.41	–0.04	–1.48:	0.15	TO/FHB	
BS 17570–0090	0.05	13.00	0.34	–0.21	–2.47	0.23	TO	
BS 17570–0092	0.03	13.92	0.48	–0.12	–0.55	0.15	TO	
BS 17570–0093	0.03	13.72	0.41	–0.13	–1.48	0.15	TO	
BS 17570–0096	0.04	12.83	0.36	–0.06	–0.87	0.15	TO	
BS 17574–0129	0.15	10.27	0.44	0.00	–2.04	0.15	FHB	
BS 17575–0052	0.09	11.03	0.29	0.10	–0.69::	0.19	FHB/A	BLUE
BS 17575–0059	0.10	10.28	0.25	–0.02	–0.44::	0.22	FHB/A	BLUE
BS 17575–0115	0.11	11.86	0.34	–0.11	–0.89	0.16	TO	
BS 17575–0128	0.26	12.68	0.27	0.09	...	...	...	NO SPEC
BS 17575–0129	0.26	10.43	0.14	0.14	...	...	...	NO SPEC
BS 17575–0142	0.18	13.16	0.28	0.04	...	...	...	NO SPEC
BS 17575–0143	0.18	11.06	0.27	0.01	...	...	...	NO SPEC
BS 17575–0144	0.17	11.51	0.19	0.04	...	...	...	NO SPEC
BS 17575–0163	0.18	12.06	0.26	0.09	–1.53::	0.25	FHB/A	BLUE
BS 17575–0168	0.13	13.00	0.42	–0.23	–2.11	0.15	TO	
BS 17575–0169	0.17	11.85	0.25	0.09	–2.02::	0.28	FHB/A	BLUE
BS 17575–0213	0.37	11.65	0.21	–0.17	–0.32::	0.22	FHB/A	BLUE
BS 17578–0001	0.11	10.40	1.09	0.97	...	...	...	CORE EM
BS 17578–0002	0.05	11.58	0.62	0.20	–0.30	0.31	TO	
BS 17578–0014	0.07	12.08	0.48	–0.07	–0.84	0.15	TO	
BS 17578–0019	0.06	12.34	0.53	–0.01	–1.18	0.20	TO	
BS 17578–0025	0.08	11.31	0.43	–0.01	–1.10:	0.15	TO/FHB	
BS 17578–0033	0.11	11.86	0.59	0.07	–1.20	0.28	SG	
BS 17578–0036	0.07	12.12	0.49	0.01	–0.84:	0.15	TO/FHB	
BS 17578–0046	0.13	11.94	0.35	0.00	–0.07	0.25	TO	
BS 17578–0051	0.11	11.59	0.44	0.02	–0.88	0.15	FHB	
BS 17578–0059	0.11	12.03	0.40	–0.09	–0.37	0.19	TO	
BS 17578–0060	0.06	11.95	0.46	–0.21	–1.45	0.15	TO	
BS 17578–0061	0.08	11.57	0.43	–0.03	–0.10	0.18	TO	
BS 17578–0062	0.12	11.16	0.39	0.16	–1.62	0.15	FHB	
BS 17578–0065	0.07	11.98	0.52	–0.14	–0.68	0.17	TO	
BS 17578–0068	0.11	12.64	0.35	–0.21	–0.31	0.21	TO	
BS 17578–0071	0.11	11.55	0.31	–0.11	–0.12	0.25	TO	
BS 17578–0087	0.11	11.63	0.26	0.01	...	...	...	NO SPEC
BS 17578–0088	0.10	12.53	0.43	–0.10	–0.59	0.15	TO	
BS 17578–0090	0.07	12.39	0.60	–0.04	–0.84	0.29	TO	

Table 5—Continued

Star (1)	$E(B - V)_A$ (2)	V_o (3)	$(B - V)_o$ (4)	$(U - B)_o$ (5)	$[\text{Fe}/\text{H}]_{\text{K3}}$ (6)	σ_{Fe} (7)	Class (8)	Comment (9)
BS 17578–0097	0.08	12.21	0.46	−0.01	−0.84:	0.15	TO/FHB	
BS 17578–0103	0.09	12.66	0.54	0.13	−1.62	0.15	TO	
BS 17578–0117	0.07	12.34	0.36	0.02	−0.46:	0.20	TO/FHB	
BS 17578–0118	0.05	11.99	0.46	−0.08	−0.77	0.15	TO	
BS 17579–0011	0.02	12.12	0.50	−0.10	−0.68	0.17	TO	
BS 17579–0016	0.02	12.36	0.41	−0.09	−0.83	0.15	TO	
BS 17579–0020	0.03	12.38	0.51	−0.14	−0.34	0.18	TO	
BS 17579–0033	0.03	12.42	0.39	−0.14	−1.21	0.15	TO	
BS 17579–0040	0.02	13.44	0.57	−0.09	...	...	...	NO SPEC
BS 17579–0047	0.02	12.38	0.52	0.00	−0.68	0.17	TO	
BS 17579–0049	0.02	12.38	0.58	−0.07	−0.84	0.29	TO	
BS 17579–0050	0.02	12.52	0.57	0.15	−0.19	0.22	TO	
BS 17579–0051	0.02	11.76	0.47	−0.10	−0.64	0.15	TO	
BS 17583–0011	0.07	11.74	0.46	−0.02	−0.50	0.15	TO	
BS 17583–0012	0.08	12.20	0.46	0.04	−0.37:	0.15	TO/FHB	
BS 17583–0013	0.08	12.09	0.54	−0.01	−1.02	0.21	TO	
BS 17583–0014	0.14	12.36	0.46	0.04	−0.50	0.15	FHB	
BS 17583–0015	0.09	11.30	0.41	0.04	−0.55:	0.15	TO/FHB	
BS 17583–0028	0.11	12.64	0.44	−0.05	−1.45	0.15	FHB	
BS 17583–0032	0.13	11.05	0.41	0.13	−1.76	0.15	FHB	
BS 17583–0057	0.05	12.89	0.53	−0.06	−0.44	0.20	TO	
BS 17583–0064	0.11	12.60	0.40	0.08	−0.07	0.19	FHB	
BS 17583–0065	0.10	12.28	0.27	0.01	−0.01::	0.26	FHB/A	BLUE
BS 17583–0067	0.11	12.22	0.28	0.21	−2.22::	0.26	FHB/A	BLUE
BS 17583–0070	0.07	12.22	0.29	0.10	−2.45::	0.23	FHB/A	BLUE
BS 17583–0077	0.05	12.20	0.46	−0.01	−0.50	0.15	TO	
BS 17583–0083	0.06	12.27	0.41	−0.25	−0.26	0.18	TO	
BS 17583–0084	0.08	12.62	0.47	0.00	−0.64:	0.15	TO/FHB	
BS 17583–0092	0.04	10.20	0.30	0.10	−0.94	0.16	TO	
BS 17583–0094	0.05	12.38	0.60	0.06	−0.64	0.28	TO	
BS 17583–0095	0.08	13.01	0.44	−0.05	−0.59	0.15	TO	
BS 17583–0097	0.06	11.94	0.65	0.13	−0.48	0.33	SG	
BS 17583–0110	0.09	10.72	0.18	0.10	−0.32::	0.23	FHB/A	BLUE
BS 17583–0116	0.04	12.11	0.62	−0.05	−0.30	0.29	TO	
BS 17583–0124	0.06	12.33	0.44	−0.04	−1.48:	0.15	TO/FHB	
BS 17583–0125	0.07	12.53	0.56	0.08	−0.85	0.25	TO	
BS 17583–0128	0.08	12.77	0.41	0.00	−0.94:	0.15	TO/FHB	
BS 17583–0132	0.05	11.92	0.58	0.05	−0.44	0.27	TO	
BS 17583–0134	0.06	11.90	0.35	−0.07	−0.29	0.24	TO	
BS 17583–0140	0.04	12.18	0.53	−0.03	−0.67	0.21	TO	
BS 17583–0146	0.05	12.05	0.44	−0.05	−1.07:	0.15	TO/FHB	
BS 17583–0152	0.05	12.05	0.52	0.04	−0.20	0.18	TO	
BS 17583–0154	0.05	12.55	0.39	−0.05	−0.47	0.17	TO	
BS 17583–0155	0.05	12.15	0.44	0.05	−0.22	0.16	TO	
BS 17583–0157	0.07	11.96	0.74	0.30	−1.20	0.38	G	
BS 17583–0160	0.05	12.75	0.56	0.14	−0.03	0.20	TO	
BS 17583–0163	0.04	12.15	0.47	0.01	−0.64:	0.15	TO/FHB	
BS 17583–0165	0.07	12.59	0.39	−0.06	−0.28	0.20	TO	
BS 17584–0005	0.04	11.66	0.50	−0.06	−0.84	0.15	TO	
BS 17584–0012	0.05	11.17	1.05	0.65	...	...	...	CORE EM
BS 17584–0016	0.04	11.92	0.61	0.05	−0.50	0.31	TO	
BS 17584–0021	0.07	12.51	0.44	−0.10	−0.59	0.15	TO	
BS 17584–0030	0.07	11.74	0.33	0.04	−0.64	0.19	FHB	
BS 17584–0032	0.06	12.74	0.51	−0.15	−1.39	0.15	TO	
BS 17584–0035	0.08	11.69	0.67	−0.04	−0.91	0.36	SG	
BS 17584–0037	0.05	12.09	0.62	0.25	−0.52	0.31	SG	
BS 17584–0038	0.05	12.13	0.65	0.03	−0.88	0.34	SG	
BS 17584–0039	0.07	12.12	0.45	−0.03	−0.34	0.15	TO	
BS 17584–0040	0.07	12.19	0.44	0.00	−0.22	0.16	TO	

Table 5—Continued

Star (1)	$E(B - V)_A$ (2)	V_o (3)	$(B - V)_o$ (4)	$(U - B)_o$ (5)	$[\text{Fe}/\text{H}]_{\text{K3}}$ (6)	σ_{Fe} (7)	Class (8)	Comment (9)
BS 17584–0041	0.07	12.36	0.51	−0.02	−0.34	0.18	TO	
BS 17584–0043	0.06	12.34	0.41	0.04	−0.49	0.15	TO	
BS 17584–0045	0.04	12.77	0.40	0.00	−0.60:	0.15	TO/FHB	
BS 17584–0046	0.05	12.59	0.31	−0.21	−0.81	0.18	TO	
BS 17584–0048	0.06	11.00	0.39	0.01	−0.75:	0.15	TO/FHB	
BS 17584–0051	0.03	9.55	0.41	−0.25	−0.83	0.15	TO	
BS 17584–0053	0.06	10.68	0.28	−0.14	−0.01::	0.26	FHB/A	BLUE
BS 17584–0061	0.05	12.68	0.45	−0.03	−1.07:	0.15	TO/FHB	
BS 17584–0070	0.03	12.26	0.49	−0.20	−1.23	0.15	TO	
BS 17584–0071	0.04	12.69	0.51	−0.05	−1.15	0.15	TO	
BS 17584–0074	0.03	9.66	0.32	−0.12	−0.57	0.21	TO	
BS 17584–0083	0.03	12.66	0.51	−0.01	−1.49	0.15	TO	
BS 17584–0084	0.04	13.09	0.46	−0.22	−0.77	0.15	TO	
BS 17584–0088	0.04	13.28	0.47	−0.24	−0.64	0.15	TO	
BS 17585–0132	0.38	11.20	0.31	0.14	−2.02	0.28	FHB	
CS 22179–0024	0.03	14.05	0.36	−0.27	−1.37:	0.19	TO	
CS 22872–0036	0.15	14.42	0.36	−0.22	−2.32	0.22	TO	
CS 22889–0045	0.04	13.19	0.42	−0.35	−1.90	0.15	TO	
CS 22962–0021	0.02	13.40	0.48	−0.20	−2.35	0.15	SG	
CS 22962–0027	0.03	14.47	0.49	−0.04	−1.51:	0.15	TO/FHB	
CS 22965–0016	0.04	13.95	0.70	0.02	−2.65	0.15	G	
CS 22965–0029	0.08	13.55	0.52	−0.08	−2.26	0.15	SG	
CS 22965–0053	0.13	12.35	0.30	0.10	−1.24	0.22	FHB	
CS 22965–0054	0.13	14.69	0.46	−0.29	−3.43	0.26	SG	
CS 29502–0001	0.06	11.21	0.36	−0.05	...	...	...	NO SPEC
CS 29502–0009	0.08	13.58	0.63	−0.12	−2.45	0.15	G	
CS 29502–0042	0.07	12.57	0.53	−0.08	−2.45	0.21	SG	
CS 29502–0062	0.04	11.18	0.46	−0.05	−0.50	0.15	TO	
CS 29502–0067	0.11	13.60	0.66	−0.10	−2.30	0.15	G	
CS 29516–0017	0.10	12.10	−0.02	0.06	...	...	...	BLUE
CS 29516–0024	0.10	13.26	0.76	0.13	−2.86	0.15	G	
CS 29516–0026	0.11	12.59	0.23	0.15	−1.38::	0.24	FHB/A	BLUE
CS 29516–0028	0.13	14.62	0.42	−0.37	−2.87	0.23	TO	
CS 29516–0032	0.10	14.23	0.56	0.01	...	...	...	CORE EM
CS 29516–0037	0.14	11.89	0.09	0.15	−3.49::	0.15	FHB/A	BLUE
CS 29516–0096	0.10	15.00	0.07	0.09	−2.45::	0.23	FHB/A	BLUE
CS 29521–0012	0.06	14.63	0.54	−0.24	−2.45	0.21	SG	
CS 29521–0031	0.06	12.62	0.55	−0.08	−1.18	0.20	TO	
CS 29521–0036	0.06	12.35	0.41	−0.09	−1.67:	0.15	TO/FHB	
CS 29521–0065	0.06	11.05	0.24	0.16	−0.57::	0.21	FHB/A	BLUE
CS 29522–0046	0.10	12.43	0.39	−0.20	−3.24	0.20	TO	
CS 30312–0006	0.14	13.21	0.48	−0.15	−2.30	0.15	SG	
CS 30312–0044	0.08	13.09	0.70	0.03	−2.59	0.15	G	
CS 30312–0067	0.13	13.83	0.44	0.02	−1.92	0.15	FHB	
CS 30325–0023	0.04	13.81	0.69	0.09	−2.09	0.16	G	
CS 30325–0033	0.03	12.85	0.41	−0.23	−1.58	0.15	TO	
CS 30332–0118	0.10	14.04	0.36	−0.23	−1.14	0.15	TO	
CS 30332–0119	0.09	15.00	0.54	0.06	−1.83	0.15	SG	
CS 30332–0138	0.10	13.67	1.08	0.76	−2.06	0.22	G	
CS 30333–0002	0.07	11.10	0.43	−0.36	−1.84	0.15	TO	
CS 31061–0004	0.04	13.43	0.56	−0.03	−1.58	0.20	SG	
CS 31061–0032	0.04	13.78	0.36	−0.31	−2.70	0.20	TO	
CS 31061–0033	0.03	13.55	0.65	−0.02	−0.61	0.37	SG	
CS 31061–0057	0.03	14.12	0.31	−0.06	−2.91	0.19	FHB	
CS 31061–0062	0.06	13.73	0.42	−0.35	−2.20	0.15	TO	
CS 31061–0072	0.04	13.85	0.44	−0.28	−1.92	0.15	TO	
CS 31061–0074	0.05	13.83	0.40	−0.27	−1.48	0.15	TO	
CS 31067–0019	0.10	12.30	0.35	−0.33	−1.42:	0.24	TO	
CS 31067–0039	0.20	13.07	0.67	0.21	−1.26	0.34	G	

Table 5—Continued

Star (1)	$E(B - V)_A$ (2)	V_o (3)	$(B - V)_o$ (4)	$(U - B)_o$ (5)	$[\text{Fe}/\text{H}]_{\text{K3}}$ (6)	σ_{Fe} (7)	Class (8)	Comment (9)
CS 31067–0045	0.11	13.51	0.58	−0.05	−0.64	0.28	TO	
CS 31069–0005	0.02	13.46	0.55	0.06	−1.03	0.25	TO	
CS 31069–0082	0.06	13.87	0.41	−0.10	−1.06	0.15	TO	
CS 31069–0109	0.05	12.75	0.82	0.46	...	...	...	CORE EM
CS 31070–0047	0.02	14.05	0.36	−0.03	−0.23	0.23	TO	
CS 31074–0005	0.18	13.66	0.49	−0.13	−0.69	0.15	TO	
CS 31079–0004	0.09	12.88	0.37	−0.23	−1.88	0.23	TO	
CS 31084–0022	0.18	11.02	0.27	−0.30	−1.53::	0.25	FHB/A	BLUE

Table 6. De-reddened Colors, Abundances, and Classifications (Variables)

Star (1)	$E(B - V)_A$ (2)	V_o (3)	$(B - V)_o$ (4)	$(U - B)_o$ (5)	$[\text{Fe}/\text{H}]_{K3}$ (6)	σ_{Fe} (7)	Class (8)	Comment (9)
BS 16083–0172	0.03	13.04	0.58	0.07	–2.37	0.15	G	
BS 16542–0002	0.06	12.43	0.70	–0.08	–2.41	0.15	G	
BS 16556–0081	0.03	14.23	0.45	–0.27	–2.24	0.15	TO	
BS 17569–0014	0.06	12.81	0.42	–0.15	–0.26	0.18	TO	
BS 17569–0023	0.05	10.90	1.39	1.35	–1.49::	0.23	G	RED
BS 17569–0027	0.05	13.83	0.44	–0.14	–1.07	0.15	TO	
BS 17569–0028	0.05	13.74	0.51	–0.20	–1.71	0.15	TO	
BS 17569–0047	0.05	14.22	0.42	–0.20	–1.97	0.15	TO	
BS 17569–0063	0.08	14.82	–0.42	–1.32	...	...	...	BLUE
BS 17569–0078	0.04	13.76	0.47	–0.19	–1.96	0.15	TO	
BS 17575–0182	0.37	11.71	0.36	0.17	–1.01	0.15	FHB	
BS 17578–0013	0.07	12.05	0.49	–0.20	–0.24	0.17	TO	
BS 17578–0032	0.10	11.80	0.37	–0.32	–0.31	0.21	TO	
BS 17578–0056	0.10	9.61	0.28	0.07	–2.02::	0.28	FHB/A	BLUE
BS 17578–0106	0.10	11.80	0.48	0.04	–0.24:	0.17	TO/FHB	
BS 17579–0012	0.02	12.22	0.08	–0.92	–2.67::	0.20	FHB/A	BLUE
BS 17583–0100	0.06	12.18	0.43	–0.24	–2.20	0.15	TO	
BS 17583–0111	0.07	12.40	0.42	–0.13	–0.49	0.15	TO	
BS 17583–0115	0.06	12.60	0.61	0.06	–0.99	0.31	SG	
BS 17584–0024	0.11	12.45	0.45	0.06	–1.15	0.15	FHB	
BS 17584–0034	0.04	13.10	0.49	–0.28	–1.23	0.15	TO	
CS 22954–0004	0.03	14.18	0.39	–0.24	–2.44	0.18	TO	
CS 30332–0012	0.09	14.37	0.45	–0.27	–2.81	0.26	SG	

Fig. 1.— (a) Comparison of the V magnitudes for 85 stars in common between the present sample, V (*our*), and that of Anthony-Twarog et al. (2000), V (*AT*). A one-to-one line is shown. (b) Comparison of the measured, $(B - V)$ (*our*), and estimated, $(B - V)_{est}(AT)$, colors for the same sample. A one-to-one line is shown (solid line), along with a locally weighted regression line (dashed line) that indicates a slight disagreement at bluer colors.

Fig. 2.— Limiting $(U - B)_0$ colors of stars in the color range $0.3 \leq (B - V)_0 \leq 0.5$, as a function of metallicity, obtained from the Revised Yale Isochrones. The region above the solid line corresponds to the locations of stars with surface gravities of main-sequence TO stars, while the region below the dashed line applies for stars with surface gravities of FHB stars. See text for additional explanation.

Fig. 3.— De-reddened two-color diagram for the non-variable stars in our sample over the color range for which reliable metallicity estimates can be obtained. A solar-abundance main sequence line, obtained from the Revised Yale Isochrones, is shown for comparison.





